

Will Elevated Carbon Dioxide Concentration Amplify the Benefits of Nitrogen Fixation in Legumes?¹

Alistair Rogers^{2*}, Elizabeth A. Ainsworth², and Andrew D.B. Leakey²

Environmental Sciences Department, Brookhaven National Laboratory, Upton, New York 11973 (A.R.); Department of Plant Biology (E.A.A., A.D.B.L.), Department of Crop Sciences (A.R., E.A.A.), and Institute for Genomic Biology (E.A.A., A.D.B.L.), University of Illinois at Urbana-Champaign, Urbana, Illinois 61801; and Photosynthesis Research Unit, United States Department of Agriculture/Agricultural Research Service, Urbana, Illinois 61801 (E.A.A.)

Growth at elevated [CO₂] stimulates photosynthesis and increases carbon (C) supply in all C₃ species. A sustained and maximal stimulation in productivity at elevated [CO₂] requires an enhanced nutrient supply to match the increase in C acquisition. The ability of legumes to exchange C for nitrogen (N) with their N₂-fixing symbionts has led to the hypothesis that legumes will have a competitive advantage over non-leguminous species when grown at elevated [CO₂]. On balance, evidence suggests that in managed systems, legumes are more responsive to elevated [CO₂] than other plants (e.g. Ainsworth and Long, 2005); however, in natural ecosystems, nutrient availability can limit the response of legumes to elevated [CO₂] (Hungate et al., 2004; van Groenigen et al., 2006). Here, we consider these observations, outline the mechanisms that underlie them, and examine recent work that advances our understanding of how legumes respond to growth at elevated [CO₂]. First we highlight the global importance of legumes and provide a brief overview of the symbiotic relationship.

IMPORTANCE OF LEGUMES

The majority of species capable of forming a symbiotic relationship with N₂-fixing bacteria belong to the Leguminosae family, and most of the roughly 20,000 legume species form a relationship with nodule-inducing bacteria, collectively known as rhizobia. The seeds of legumes have a high protein content, and some are also rich in oil. Consequently, legumes are an important source of protein and calories for humans and animals, exemplified by the fact that 25 countries plant 20% to 57% of their arable land with legumes

such as Bambara groundnuts (*Vigna subterranea*), peas (*Pisum sativum*), beans (*Phaseolus vulgaris*), and lentils (*Lens culinaris*; <http://faostat.fao.org/>; Fig. 1). These include countries using intensive farming practices that produce mainly soybean (*Glycine max*), as well as developing countries where smallholder and subsistence farmers grow a wide variety of species that contribute to local food security. In addition, legumes are important components of animal forage in both temperate and tropical zones (Kretschmer and Pitman, 2001). In the tropics leguminous trees are used in rotation, or in multispecies cropping systems to enhance soil fertility, and as a source of wood. In all row crop, pasture, and agroforestry systems, the ability of legumes to fix N₂ reduces the need for chemical fertilizer inputs and provides important economic and environmental benefits (Peoples et al., 1995). Between one-third and one-half of the total N added to agricultural land is attributable to the legume-rhizobia symbiosis (Herridge et al., 2008; <http://faostat.fao.org/>). The contribution of legume-rhizobia symbioses to N₂-fixation natural ecosystems is hard to estimate, but is important to the composition and function of many terrestrial ecosystems, with woody legumes being particularly abundant in the tropics, while herbaceous legumes are widespread at higher latitudes (Vitousek et al., 2002).

THE SYMBIOSIS

The formation of a root nodule is preceded by an elaborate molecular courtship and the controlled infection of the plant host cell by rhizobia. Within the root nodule, a bacteroid, containing differentiated rhizobia, is isolated from the host cell by a peribacteroid membrane. In addition to forming and maintaining the root nodule, the host provides the bacteroid with a supply of C to fuel N₂ fixation. Suc is delivered to the root nodule via the phloem where it is cleaved by Suc synthase, enters glycolysis, and ultimately provides malate to the bacteroid. The bacteroid returns NH₄⁺ to the host cell, which is assimilated into Gln. Further metabolism into other N-containing transport compounds uses C skeletons furnished by the host, and in the case of N-rich ureides, takes place in adjacent uninfected host cells. Both amides (Gln and

¹ This work was supported in part by the U.S. Department of Energy Office of Science (grant no. DE-AC02-98CH10886 to A.R. and Brookhaven National Laboratory).

² These authors contributed equally to the article.

* Corresponding author; e-mail arogers@bnl.gov.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Alistair Rogers (arogers@bnl.gov).

www.plantphysiol.org/cgi/doi/10.1104/pp.109.144113




Figure 1. Percentage of agricultural land per country used for production of leguminous food crops in 2006. National crop production data were taken from the ProdSTAT database, Food and Agriculture Organization of the United Nations (<http://faostat.fao.org>), and included the following major crops: Bambara groundnuts, broad beans (*Vicia faba*), carob (*Ceratonia siliqua*), cowpeas (*Vigna unguiculata*), chickpeas (*Cicer arietinum*), dry beans (*Phaseolus* spp. and *Vigna* spp.), lentils, peanut, peas, pigeon peas (*Cajanus cajan*), and soybean.

Asn) and uriedes are exported from the root nodule via the xylem (Fig. 2). Within the bacteroid, malate is oxidized by the Krebs cycle to provide reductant for the nitrogenase complex and for the respiratory chain that fuels the nitrogenase complex with ATP. The outer layer of the root nodule forms a diffusive barrier that limits the entry of O₂ and protects the highly sensitive nitrogenase complex from O₂ inhibition. However, this barrier also limits the supply of O₂ to the bacteroid respiratory chain. The presence of high concentrations of leghemoglobin in the host cell, which gives active nodules their characteristic pink color, increases the flux of O₂ through the cytoplasm and controls the concentration of free O₂ available to the respiratory chain in the peribacteroid membrane (Crawford et al., 2000; Fig. 2).

ARE LEGUMES MORE RESPONSIVE TO ELEVATED [CO₂] THAN OTHER PLANTS?

The hypothesis that legumes have an advantage over nonleguminous plants in elevated [CO₂] is prevalent in the literature (e.g. Soussana and Hartwig, 1996; Zanetti et al., 1996; Serraj et al., 1998; Rogers et al., 2006). This hypothesis is well supported by numerous studies that report greater stimulation of photosynthesis and growth of legume species compared to other functional groups (e.g. Reich et al., 2001; Lee et al., 2003; Ainsworth and Long, 2005). Furthermore, experiments comparing nodulating and non-nodulating genotypes within a legume species have found that N fixers are more responsive to elevated [CO₂] than nonfixers (Nakamura et al., 1999; Luscher et al., 2000; Ainsworth et al., 2002, 2004). However, there is also significant variation in the magnitude of the response to elevated [CO₂] among legume species and genotypes (Reich et al., 2001; West et al., 2005), and

legume aboveground primary production was not statistically greater than that of other functional types when averaged across four Free Air CO₂ Enrichment (FACE) experiments (Nowak et al., 2004). A recent meta-analysis suggested that elevated [CO₂] will have no effect on N₂ fixation in many natural terrestrial ecosystems, which are characterized by intact soils, naturally occurring plant communities, and no fertilizer amendments (van Groenigen et al., 2006).

WHY IS ELEVATED [CO₂] PARTICULARLY BENEFICIAL FOR LEGUMES? THE UNDERLYING MECHANISMS

In C₃ plants, elevated [CO₂] increases photosynthesis and decreases stomatal conductance. All other effects at the plant and ecosystem level stem from these two primary responses and the secondary effects they elicit (Ainsworth and Rogers, 2007). In the absence of other nutrient limitations, legumes will be able to capitalize on the benefits of elevated [CO₂] by both increasing N₂ fixation and by reducing the negative impact of drought on N₂ fixation. In addition, legumes will be able to avoid or limit the few deleterious effects of elevated [CO₂] by reducing the potential for sink limitation of C acquisition and limiting the reductions in leaf and grain N content that are typically observed in C₃ plants grown at elevated [CO₂]. Here, we examine the mechanisms that allow legumes




Figure 2. A root nodule showing the exchange of C and N between the host and bacteroid. Leg, Leghemoglobin; RC, bacterial respiratory chain; NC, the nitrogenase complex; SS, Suc synthase.

to both capitalize on the benefits, and avoid, or limit, the disadvantages of growth at elevated [CO₂].

DO LEGUMES INCREASE N₂ FIXATION AT ELEVATED [CO₂]?

In fertilized crops and pasture species, the hypothesis that greater photoassimilate availability at elevated CO₂ supports stimulated N₂ fixation is broadly supported (e.g. Soussana and Hartwig, 1996; Zanetti et al., 1996; Ross et al., 2004; Rogers et al., 2006; Fig. 3). FACE experiments with soybean (*Glycine max*) and clover (*Trifolium repens*) showed a stimulation in photosynthetic C uptake and leaf carbohydrate content at elevated [CO₂] (Ainsworth et al., 2003b; Rogers et al., 2004; Leakey et al., 2009; Fig. 3). Increases in soybean aboveground biomass were not accompanied by reductions in tissue N content, implying that increased C supply was matched by additional N₂ fixation at elevated [CO₂] (Morgan et al., 2005; Rogers et al., 2006). An increase in the relative contribution of symbiotically fixed N to total plant N at elevated [CO₂] has been reported in a number of experiments (e.g. Schortemeyer et al., 2002; Lee et al., 2003), and in clover, the increased relative contribution of symbiotically fixed N₂ at elevated [CO₂] was a critical factor in maintaining the C to N ratio in the ecosystem as a whole (Zanetti et al., 1996). These experiments provide ample evidence that legumes increase N₂ fixation at elevated [CO₂].

An increase in N₂ fixation at elevated [CO₂] could result from an increase in nodule size, nodule number

per plant, or an increase in specific nitrogenase activity, i.e. the amount of N₂ fixed per unit nodule and time (Fig. 3). Increased nodule biomass per plant is commonly reported in leguminous trees (e.g. Schortemeyer et al., 2002) and herbaceous species (e.g. Lee et al., 2003) grown at elevated [CO₂]. Changes in both nodule size and nodule number contribute to increased nodule biomass in different species and studies. On the other hand, there is less evidence that specific nitrogenase activity increases at elevated [CO₂] (Cen and Layzell, 2004), perhaps because many legumes lack excess specific nitrogenase capacity (Vance and Heichel, 1991).

In contrast to N₂ fixation in fertilized agricultural environments, N₂ fixation in natural ecosystems is not consistently stimulated by elevated [CO₂] (van Groenigen et al., 2006). Why? In some cases, stimulation of N₂ fixation by elevated [CO₂] may be species specific (West et al., 2005) or driven by the effects of elevated [CO₂] on competition and abundance of legumes in the ecosystem (Reich et al., 2006). In other cases, stimulation of N₂ fixation by elevated [CO₂] is limited by the availability of other essential elements (Reich et al., 2006; van Groenigen et al., 2006). For example, in a 6-year study of a N-poor calcareous grassland, phosphorus (P) limited the response of legumes to elevated [CO₂], which in turn limited N transfer to nonleguminous species in the ecosystem (Niklaus and Körner, 2004). A study of a Florida scrub-oak ecosystem reported that the initial stimulation of N₂ fixation in *Galactia elliotii* disappeared over time as molybdenum (Mo) became limiting in elevated [CO₂] (Hungate et al., 2004). Mo is a required cofactor for the




Figure 3. Summary of the effects of elevated [CO₂] and nutrient supply on legume leaf, pod/seed, and nodule characteristics. g_s , Stomatal conductance; $V_{c,max}$, maximum photosynthetic carboxylation capacity; J_{max} , maximum capacity for electron transport leading to ribulose-1,5-bisP regeneration. + indicates increases in elevated [CO₂]; - indicates decreases in elevated [CO₂]. The number of + or - symbols indicates the magnitude of change at elevated [CO₂].

nitrogenase complex, and elevated $[\text{CO}_2]$ was thought to increase Mo adsorption to soil particles, decreasing the availability of Mo, and causing a systematic decline in N_2 fixation (Hungate et al., 2004).

These apparent nutrient limitations of N_2 fixation at elevated $[\text{CO}_2]$ may not be specific to legumes or physiologically distinct from nutrient limitation in nonleguminous species. Mo limitation, which is largely confined to acidic soils, may impact N_2 fixation directly (Hungate et al., 2004). However, Mo limitation is also likely to affect plants dependent on soilborne nitrate because nitrate reductase also requires Mo cofactors (Crawford et al., 2000). Consider the P limitation of N_2 fixation; less than 1% of total plant P is found in nodules, and plants grown with a low P availability often have higher specific N_2 fixation (e.g. Almeida et al., 2000). Some crop species do need more P when they acquire N through N_2 fixation. However, in wild species, where the nutrient limitation of N_2 fixation has been shown to constrain the response to elevated $[\text{CO}_2]$ (van Groenigen et al., 2006), the P requirement of legumes fixing N_2 is typically the same as when they acquired N from a mineral source (Vitousek et al., 2002). These observations support the view that the process of N_2 fixation is not limited directly by P availability and that whole-plant P requirement does not increase with N_2 fixation. Therefore, the observation that N_2 fixation, and the positive growth response of legumes, is constrained by a limiting P supply at elevated $[\text{CO}_2]$ probably reflects a general P limitation that is exacerbated at elevated $[\text{CO}_2]$. The work conducted at the Swiss FACE experiment illustrates this well (Almeida et al., 2000). When clover was grown at a limiting P supply, nodule mass was decreased but specific N_2 fixation was higher. These plants also had a higher leaf N content, and a higher whole-plant N to P ratio than P-sufficient plants. When these plants were grown at elevated $[\text{CO}_2]$, leaf starch content was increased. This indicates that plants grown at elevated $[\text{CO}_2]$ and a limiting P supply were not limited by C or N supply. Rather, growth in low P limited the demand for N, resulting in fewer nodules and lower total N_2 fixation (Almeida et al., 2000). When P availability was increased, growth at elevated $[\text{CO}_2]$ resulted in increased plant growth. The internal mechanism of response to limiting P under elevated $[\text{CO}_2]$ is probably not unique to legumes. However, they may be unusual in the extent to which growth at elevated $[\text{CO}_2]$ may help plants to release P from normally unavailable sources. The additional C and N_2 fixed at elevated $[\text{CO}_2]$ by legumes can be partitioned to promote root growth, fuel mycorrhizal associations, increase the secretion of chelating agents, and increase the secretion of extracellular phosphatases (Vitousek et al., 2002; Houlton et al., 2008).

Therefore, the non-N nutrient limitation of the response of legumes to elevated $[\text{CO}_2]$ is likely a general nutrient limitation of plant growth that is exacerbated at elevated $[\text{CO}_2]$ where the availability of C, N, and

potentially water is improved, rather than a specific nutrient requirement of legumes grown at elevated $[\text{CO}_2]$.

ARE LEAF AND GRAIN QUALITY ALTERED BY GROWTH AT ELEVATED $[\text{CO}_2]$ IN LEGUMES?

Reductions in leaf N content and or increases in C to N ratio are commonly observed in plants grown at elevated $[\text{CO}_2]$ (Ainsworth and Long, 2005). Reduced leaf N content negatively impacts forage quality and can result in additional feeding by protein-limited insects as they attempt to compensate for the reduced leaf N content (Whittaker, 1999). In addition, a lower leaf N content, relative to the enhanced C supply at elevated $[\text{CO}_2]$, limits the supply of N available for translocation to reproductive tissue during grain fill.

Evidence from mixed semigrassland communities managed as a forage crop showed that leaf N content and C to N ratio were unaffected by growth at elevated $[\text{CO}_2]$ in legumes, but decreased in nonlegumes (Winkler and Herbst, 2004). In field-grown soybean, Rogers et al. (2006) showed that early season reductions in leaf N content at elevated $[\text{CO}_2]$ disappeared once N_2 fixation was under way. These data provide good evidence that legumes that lack restrictions on their symbiotic exchange will not exhibit a reduced leaf N content at elevated $[\text{CO}_2]$, and that the quality of leguminous forage crops will not be reduced (Fig. 3). There is evidence that this also helps legumes avoid compensatory feeding at elevated $[\text{CO}_2]$ (Karowe, 2007; Lau et al., 2008). However, additional folivory at elevated $[\text{CO}_2]$ was reported in soybean, and the response was linked to increased sugar content and a compromised defense system. This indicates that despite avoiding N dilution, increased damage by herbivores will likely occur in some legume species (Hamilton et al., 2005; Zavala et al., 2008, 2009). The response of herbivory to rising $[\text{CO}_2]$ is highly complex and at this time a clear consensus of the response of herbivores to legumes grown at elevated $[\text{CO}_2]$ is unclear.

Elevated $[\text{CO}_2]$ will have direct effects on human health and well being through changes in grain quality. Protein content of nonleguminous C_3 grains including barley (*Hordeum vulgare*), wheat (*Triticum aestivum*), and rice (*Oryza sativa*) declined by 14% at elevated $[\text{CO}_2]$ (Taub et al., 2008), and concentrations of calcium, sulfur, iron, and zinc were also significantly lower in wheat grain (Loladze, 2002). Will elevated $[\text{CO}_2]$ affect legume seed quality to a similar extent? The evidence reported in two meta-analyses suggests that seed protein and N content will not be as affected by elevated $[\text{CO}_2]$ in legumes as nonlegumes (Jablonski et al., 2002; Taub et al., 2008; Fig. 3). In soybean, growth at elevated $[\text{CO}_2]$ significantly decreased protein content, but only by 1.4% (Taub et al., 2008). Averaging the effects of elevated $[\text{CO}_2]$ on wild and domestic legumes, there was no effect on seed N

(Jablonski et al., 2002). Other important aspects of seed quality are also maintained at elevated [CO₂] in legumes. For example, percent oil in seeds of red kidney bean (*Phaseolus vulgaris*), peanut (*Arachis hypogaea*), mung bean (*Vigna mungo*), and soybean were unaffected by elevated [CO₂] (Burkey et al., 2007; Ziska et al., 2007; Thomas et al., 2009). The seed oil composition was altered by elevated [CO₂] in some species and experiments, but in general, results do not cause concern for rising [CO₂] effects on composition and edible quality of legumes (e.g. Burkey et al., 2007). Legumes can also provide key minerals, including iron, copper, manganese, calcium, magnesium, zinc, cobalt, P, and potassium, but there is very little data available to describe how mineral content of legume seeds changes at elevated [CO₂]. Theoretically, those minerals that are delivered to the plant through the transpiration stream, including P, may be less available at elevated [CO₂] but this phenomena would not be unique to legumes.

WILL LEGUMES MAXIMIZE PHOTOSYNTHETIC STIMULATION BY ELEVATED [CO₂]?

The sustained stimulation of C₃ photosynthesis by elevated [CO₂] increases the availability of carbohydrate and in doing so alters the balance of C and N resources. As a result the plant acclimates, adjusting its photosynthetic, respiratory, and N metabolism to improve performance under elevated [CO₂] (Leakey et al., 2009). The acclimation of photosynthesis is best understood in the context of the relationship between changes in intercellular [CO₂] (c_i) and the light-saturated rate of photosynthesis (A), otherwise known as the A - c_i curve (Fig. 4). Differences in the impact of elevated [CO₂] on photosynthesis among major plant functional types can be explained on the basis of variation in the A - c_i relationship (Ainsworth and Rogers, 2007). The distinct effects of elevated [CO₂] on leguminous and nonleguminous C₃ herbs can be illustrated by comparing the responses of nonleguminous perennial ryegrass (*Lolium perenne*) and leguminous clover (Fig. 4). Perennial ryegrass grown at elevated [CO₂] displayed photosynthetic acclimation with a lower maximum photosynthetic carboxylation capacity ($V_{c,max}$) and a lower maximum capacity for electron transport leading to ribulose-1,5-bisP regeneration (J_{max}) when compared with plants grown at current [CO₂] (Ainsworth et al., 2003a). This results in a less-steep initial slope and lower asymptote on the A - c_i curve at elevated [CO₂] compared to current [CO₂] (Fig. 4A). As a result, leaves measured at an atmospheric [CO₂] of 600 $\mu\text{mol mol}^{-1}$ (corresponding to a c_i of approximately 420 $\mu\text{mol mol}^{-1}$) had lower A if grown at elevated [CO₂] (Elev₆₀₀ in Fig. 4A) compared to plants grown at current [CO₂] (Curr₆₀₀ in Fig. 4A). As a consequence, the stimulation of A at an atmospheric [CO₂] of 600 $\mu\text{mol mol}^{-1}$ relative to 360 $\mu\text{mol mol}^{-1}$ (corresponding to a c_i of approximately 250 $\mu\text{mol mol}^{-1}$) was less in plants grown at elevated [CO₂]




Figure 4. The response of photosynthesis (A) to internal [CO₂] (c_i) in perennial ryegrass (A) and clover (B) grown at current [CO₂] (approximately 360 $\mu\text{mol mol}^{-1}$, solid lines) and elevated [CO₂] (approximately 600 $\mu\text{mol mol}^{-1}$, broken lines) as part of the Swiss FACE experiment. Arrows indicate A at the c_i when leaves were: grown and measured at current [CO₂] (Curr₃₆₀), grown at current [CO₂] and measured at elevated [CO₂] (Curr₆₀₀), or grown and measured at elevated [CO₂] (Elev₆₀₀).

compared to plants grown at current [CO₂]. In contrast, clover did not photosynthetically acclimate to elevated [CO₂], maintaining the same $V_{c,max}$ and J_{max} at current and elevated [CO₂] and maximizing the stimulation of A by elevated [CO₂] (Fig. 4B; Ainsworth et al., 2003b). Legumes, such as clover, maximize the stimulation of productivity at elevated [CO₂] by avoiding photosynthetic acclimation because of their ability to reset the balance of C and N metabolism at elevated [CO₂]. Legumes avoid acclimation by allocating some of the additional photosynthate produced at elevated [CO₂] to their symbiosis with rhizobia to stimulate N₂ fixation. In nonleguminous species where N acquisition cannot be stimulated in this manner, C and N metabolism are balanced by reducing synthesis of Rubisco, which leads to lower $V_{c,max}$ and A , but also decreases the photosynthetic demand for the limiting nutrient, N, making it available to other sinks (Leakey et al., 2009). These

mechanisms maximize the efficiency of resource use for both functional groups, but explain how legumes can gain greater benefit from elevated $[\text{CO}_2]$ than nonlegumes.

The capacity for legumes to coordinate enhanced assimilation of C and N at elevated $[\text{CO}_2]$ to avoid down-regulation of photosynthetic capacity and ultimately maximize gains in productivity has been demonstrated across a range of species, environmental conditions, and ecological settings (Ainsworth and Rogers, 2007; Leakey et al., 2009). Nevertheless, there are situations where N_2 fixation cannot be stimulated in response to greater photosynthate availability at elevated $[\text{CO}_2]$ because some other factor is limiting. These include the nutrient limitations discussed above, as well as low temperature (Ainsworth et al., 2003b) and potentially drought stress. Unlike the response during warm spring weather (Fig. 4), low temperature was observed to cause photosynthetic acclimation to elevated $[\text{CO}_2]$ in clover (Ainsworth et al., 2003b). It seems likely that when some environmental factor prevents stimulation of N_2 fixation at elevated $[\text{CO}_2]$, excess photosynthate is likely to accumulate and lead to photosynthetic acclimation, but direct evidence of this mechanism in operation is needed.

WILL ELEVATED $[\text{CO}_2]$ HELP LEGUMES AVOID DROUGHT-INDUCED REDUCTIONS IN N_2 FIXATION?

Drought is a major limitation to crop production and causes a marked inhibition of N_2 fixation in legumes. This is particularly important in arid regions where legumes are also major food crops (Fig. 1). In some species, N_2 fixation is especially sensitive to even modest soil drying and reductions in N_2 fixation are observed before effects on transpiration or photosynthesis are apparent (Serraj et al., 1999). Despite considerable attention, the causes of drought-induced inhibition of N_2 fixation are still not well understood. There are a number of mechanisms proposed to account for the reduction in N_2 fixation under drought stress: (1) oxygen limitation, (2) C shortage, (3) regulation by N metabolism, and (4) oxidative damage of cellular components in the nodules (Serraj et al., 1999; Galvez et al., 2005; Ladrera et al., 2007; Marino et al., 2007; Naya et al., 2007). A detailed review of these mechanisms is beyond the scope of this *Update*. However, given that elevated $[\text{CO}_2]$ has previously been reported to ameliorate drought-induced reductions in N_2 fixation through a number of these mechanisms, we consider recent work in this area.

The supply of C to the bacteroid can limit N_2 fixation. Plants growing at elevated $[\text{CO}_2]$ should be able to enhance C supply to the nodule through increased photosynthesis. However, the drought-induced C shortage in the bacteroid occurs downstream of Suc supply. It is correlated with a crash in

host cell Suc synthase activity (Fig. 2), an accumulation of Suc, and a decline in malate content (Galvez et al., 2005). Therefore, it is unlikely that additional Suc available to host cell in plants grown at elevated $[\text{CO}_2]$ will be able to ameliorate a drought-induced reduction in N_2 fixation. It is not clear if this mechanism is operating in all legume species and new evidence suggests that under moderate drought, impaired respiratory activity, and oxidative damage of nodule components precedes reductions in Suc synthase activity (Naya et al., 2007; Sassi et al., 2008).

Drought-associated decreases in N_2 fixation have been linked to increases in ureides, amides, and other amino acids in the leaf and nodule, and there is evidence for both a systemic and local feedback on N_2 fixation that is associated with N metabolites (Serraj et al., 1999). Growth at elevated $[\text{CO}_2]$ would provide more C skeletons to metabolize foliar ureides and amides, and the increased N sink at elevated $[\text{CO}_2]$ would stimulate ureide catabolism and reduce the leaf N-derived signal. However, recent work has shown that control is at the nodule level and does not appear to be mediated through a systemic N response (Ladrera et al., 2007; Marino et al., 2007). Since the availability of C skeletons required for the synthesis of N transport compounds is also impacted by the drought associated crash in Suc synthase activity, there is little potential for elevated $[\text{CO}_2]$ to increase C supply and alleviate drought-induced N metabolite accumulation in the nodules.

Elevated $[\text{CO}_2]$ perhaps has the most potential to help protect N_2 fixation from moderate drought by maintaining higher soil moisture content. There is overwhelming evidence that growth at elevated $[\text{CO}_2]$ reduces stomatal conductance (g_s ; Fig. 3), and that this instantaneous response of stomata to CO_2 is maintained with time (Leakey et al., 2009). Recent evidence from FACE experiments, where the natural coupling between plants and the atmosphere is largely unaltered, also suggests that the decrease in g_s at elevated $[\text{CO}_2]$ often translates to a proportional reduction in canopy transpiration and an increased soil moisture content in a range of plants, including soybean (Leakey et al., 2009). Therefore, regardless of the mechanism by which drought impacts N_2 fixation, reduced g_s at elevated $[\text{CO}_2]$ may help prevent or delay reductions in N_2 fixation associated with dry periods by maintaining soil moisture content above inhibition thresholds. This will be particularly important for moderate drought stress where inhibition is readily reversed upon rewetting. Given that the reduction in g_s at elevated $[\text{CO}_2]$ is conserved across functional groups, legumes in mixed communities will also benefit from the increased water use efficiency of their competitors.

CONCLUSION

Current evidence suggests there are three key features of the response of legumes to elevated $[\text{CO}_2]$: (1)

unlike other nonleguminous C₃ plants, only legumes have the potential to maximize the benefit of elevated [CO₂] by matching stimulated photosynthesis with increased N₂ fixation; (2) this potential can only be realized in the absence of limitations on productivity such as nutrient deficiency, low temperature, or drought; (3) rising [CO₂] may offer some protection from drought-induced decreases in N₂ fixation, which will become more prevalent with projected changes in precipitation intensity and frequency that are projected to accompany the rise in [CO₂]. However, despite the considerable importance of legumes to both agriculture and the function of natural ecosystems, there are still key knowledge gaps. There have been very few long-term studies of the response of field-grown legumes to elevated [CO₂]. This greatly limits characterization of the environmental conditions under which N₂ fixation can or cannot be stimulated at elevated [CO₂]. The feedback effects of nutrient limitation on N₂ fixation and photosynthesis have not been quantified. Only a single leguminous food crop (soybean) has been the subject of a fully open-air CO₂ enrichment experiment, and this study has not yet reported the effects of elevated [CO₂] on N₂ fixation. No study we are aware of has quantitatively assessed the flow of C to nodules at current and elevated [CO₂]. These and other challenges create the prospect of many new and exciting findings in this subject area.

ACKNOWLEDGMENTS

We thank Joe Nivens for the soybean illustration and Craig Yendrek for comments on an earlier version of the manuscript.

Received July 2, 2009; accepted September 2, 2009; published September 15, 2009.

LITERATURE CITED

- Ainsworth EA, Davey PA, Bernacchi CJ, Dermody OC, Heaton EA, Moore DJ, Morgan PB, Naidu SL, Ra HSY, Zhu XG, et al (2002) A meta-analysis of elevated [CO₂] effects on soybean (*Glycine max*) physiology, growth and yield. *Glob Change Biol* 8: 695–709
- Ainsworth EA, Davey PA, Hymus GJ, Osborne CP, Rogers A, Blum H, Nosberger J, Long SP (2003a) Is stimulation of leaf photosynthesis by elevated carbon dioxide concentration maintained in the long term? A test with *Lolium perenne* grown for 10 years at two nitrogen fertilization levels under Free Air CO₂ Enrichment (FACE). *Plant Cell Environ* 26: 705–714
- Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy. *New Phytol* 165: 351–371
- Ainsworth EA, Rogers A (2007) The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions. *Plant Cell Environ* 30: 258–270
- Ainsworth EA, Rogers A, Blum H, Nosberger J, Long SP (2003b) Variation in acclimation of photosynthesis in *Trifolium repens* after eight years of exposure to Free Air CO₂ Enrichment (FACE). *J Exp Bot* 54: 2769–2774
- Ainsworth EA, Rogers A, Nelson R, Long SP (2004) Testing the “source-sink” hypothesis of down-regulation of photosynthesis in elevated [CO₂] in the field with single gene substitutions in *Glycine max*. *Agric Meteorol* 122: 85–94
- Almeida JPF, Hartwig UA, Frehner M, Nosberger J, Luscher A (2000) Evidence that P deficiency induces N feedback regulation of symbiotic N-2 fixation in white clover (*Trifolium repens* L.). *J Exp Bot* 51: 1289–1297
- Burkey KO, Booker FL, Pursley WA, Heagle AS (2007) Elevated carbon dioxide and ozone effects on peanut: II. Seed yield and quality. *Crop Sci* 47: 1488–1497
- Cen YP, Layzell DB (2004) Does oxygen limit nitrogenase activity in soybean exposed to elevated CO₂? *Plant Cell Environ* 27: 1229–1238
- Crawford NM, Kahn ML, Leustek T, Long SR (2000) Nitrogen and sulfur. In BB Buchanan, W Gruissem, RL Jones, eds, *Biochemistry & Molecular Biology of Plants*. American Society of Plant Physiologists, Rockville, MD, pp 786–849
- Galvez L, Gonzalez EM, Arrese-Igor C (2005) Evidence for carbon flux shortage and strong carbon/nitrogen interactions in pea nodules at early stages of water stress. *J Exp Bot* 56: 2551–2561
- Hamilton JG, Dermody O, Aldea M, Zangerl AR, Rogers A, Berenbaum MR, DeLucia EH (2005) Anthropogenic changes in tropospheric composition increase susceptibility of soybean to insect herbivory. *Environ Entomol* 34: 479–485
- Herridge DF, Peoples MB, Boddey RM (2008) Global inputs of biological nitrogen fixation in agricultural systems. *Plant Soil* 311: 1–18
- Houlton BZ, Wang YP, Vitousek PM, Field CB (2008) A unifying framework for dinitrogen fixation in the terrestrial biosphere. *Nature* 454: 327–334
- Hungate BA, Stiling PD, Dijkstra P, Johnson DW, Ketterer ME, Hymus GJ, Hinkle CR, Drake BG (2004) CO₂ elicits long-term decline in nitrogen fixation. *Science* 304: 1291
- Jablonski LM, Wang XZ, Curtis PS (2002) Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports on 79 crop and wild species. *New Phytol* 156: 9–26
- Karowe DN (2007) Are legume-feeding herbivores buffered against direct effects of elevated carbon dioxide on host plants? A test with the sulfur butterfly, *Colias philodice*. *Glob Change Biol* 13: 2045–2051
- Kretschmer AE, Pitman WD (2001) Germplasm resources of tropical forage legumes. In A Sotomayor-Rios, WD Pitman, eds, *Tropical Forage Plants*. CRC Press, Boca Raton, FL, pp 41–52
- Ladrera R, Marino D, Larrainzar E, Gonzalez EM, Arrese-Igor C (2007) Reduced carbon availability to bacteroids and elevated ureides in nodules, but not in shoots, are involved in the nitrogen fixation response to early drought in soybean. *Plant Physiol* 145: 539–546
- Lau JA, Strengbom J, Stone LR, Reich PB, Tiffin P (2008) Direct and indirect effects of CO₂, nitrogen, and community diversity on plant-enemy interactions. *Ecology* 89: 226–236
- Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR (2009) Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *J Exp Bot* 60: 2859–2876
- Lee TD, Tjoelker MG, Reich PB, Russelle MP (2003) Contrasting growth response of an N-2-fixing and non-fixing forb to elevated CO₂: dependence on soil N supply. *Plant Soil* 255: 475–486
- Loladze I (2002) Rising atmospheric CO₂ and human nutrition: toward globally imbalanced plant stoichiometry? *Trends Ecol Evol* 17: 457–461
- Luscher A, Hartwig UA, Suter D, Nosberger J (2000) Direct evidence that symbiotic N-2 fixation in fertile grassland is an important trait for a strong response of plants to elevated atmospheric CO₂. *Glob Change Biol* 6: 655–662
- Marino D, Frendo P, Ladrera R, Zabalza A, Puppo A, Arrese-Igor C, Gonzalez EM (2007) Nitrogen fixation control under drought stress: localized or systemic? *Plant Physiol* 143: 1968–1974
- Morgan PB, Bollero GA, Nelson RL, Dohleman FG, Long SP (2005) Smaller than predicted increase in aboveground net primary production and yield of field-grown soybean under fully open-air [CO₂] elevation. *Glob Change Biol* 11: 1856–1865
- Nakamura T, Koike T, Lei T, Ohashi K, Shinano T, Tadano T (1999) The effect of CO₂ enrichment on the growth of nodulated and non-nodulated isogenic types of soybean raised under two nitrogen concentrations. *Photosynthetica* 37: 61–70
- Naya L, Ladrera R, Ramos J, Gonzalez EM, Arrese-Igor C, Minchin FR, Becana M (2007) The response of carbon metabolism and antioxidant defenses of alfalfa nodules to drought stress and to the subsequent recovery of plants. *Plant Physiol* 144: 1104–1114
- Niklaus PA, Körner C (2004) Synthesis of a six-year study of calcareous grassland responses to in situ CO₂ enrichment. *Ecol Monogr* 74: 491–511
- Nowak RS, Ellsworth DS, Smith SD (2004) Functional responses of plants to elevated atmospheric CO₂—do photosynthetic and productivity data

- from FACE experiments support early predictions? *New Phytol* **162**: 253–280
- Peoples MB, Herridge DF, Ladha JK** (1995) Biological nitrogen-fixation—an efficient source of nitrogen for sustainable agricultural production. *Plant Soil* **174**: 3–28
- Reich PB, Hungate BA, Luo YQ** (2006) Carbon-nitrogen interactions in terrestrial ecosystems in response to rising atmospheric carbon dioxide. *Annu Rev Ecol Evol Syst* **37**: 611–636
- Reich PB, Tilman D, Craine J, Ellsworth D, Tjoelker MG, Knops J, Wedin D, Naeem S, Bahaeddin D, Goth J, et al** (2001) Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO₂ and N availability regimes? A field test with 16 grassland species. *New Phytol* **150**: 435–448
- Rogers A, Allen DJ, Davey PA, Morgan PB, Ainsworth EA, Bernacchi CJ, Cornic G, Dermody O, Heaton EA, Mahoney J, et al** (2004) Leaf photosynthesis and carbohydrate dynamics of soybeans grown throughout their life-cycle under Free-Air Carbon dioxide Enrichment. *Plant Cell Environ* **27**: 449–458
- Rogers A, Gibon Y, Stitt M, Morgan PB, Bernacchi CJ, Ort DR, Long SP** (2006) Increased C availability at elevated carbon dioxide concentration improves N assimilation in a legume. *Plant Cell Environ* **29**: 1651–1658
- Ross DJ, Newton PCD, Tate KR** (2004) Elevated [CO₂] effects on herbage production and soil carbon and nitrogen pools and mineralization in a species-rich, grazed pasture on a seasonally dry sand. *Plant Soil* **260**: 183–196
- Sassi S, Gonzalez EM, Aydi S, Arrese-Igor C, Abdely C** (2008) Tolerance of common bean to long-term osmotic stress is related to nodule carbon flux and antioxidant defenses: evidence from two cultivars with contrasting tolerance. *Plant Soil* **312**: 39–48
- Schortemeyer M, Atkin OK, McFarlane N, Evans JR** (2002) N₂ fixation by *Acacia* species increases under elevated atmospheric CO₂. *Plant Cell Environ* **25**: 567–579
- Serraj R, Sinclair TR, Allen LH** (1998) Soybean nodulation and N₂ fixation response to drought under carbon dioxide enrichment. *Plant Cell Environ* **21**: 491–500
- Serraj R, Sinclair TR, Purcell LC** (1999) Symbiotic N₂ fixation response to drought. *J Exp Bot* **50**: 143–155
- Soussana JF, Hartwig UA** (1996) The effects of elevated CO₂ on symbiotic N₂ fixation: a link between the carbon and nitrogen cycles in grassland ecosystems. *Plant Soil* **187**: 321–332
- Taub DR, Miller B, Allen H** (2008) Effects of elevated CO₂ on the protein concentration of food crops: a meta-analysis. *Glob Change Biol* **14**: 565–575
- Thomas JMG, Prasad PVV, Boote KJ, Allen LH** (2009) Seed composition, seedling emergence and early seedling vigour of red kidney bean seed produced at elevated temperature and carbon dioxide. *J Agron Crop Sci* **195**: 148–156
- van Groenigen KJ, Six J, Hungate BA, de Graaff MA, van Breemen N, van Kessel C** (2006) Element interactions limit soil carbon storage. *Proc Natl Acad Sci USA* **103**: 6571–6574
- Vance CP, Heichel GH** (1991) Carbon in N₂ fixation—limitation or exquisite adaptation. *Annu Rev Plant Physiol Plant Mol Biol* **42**: 373–392
- Vitousek PM, Cassman K, Cleveland C, Crews T, Field CB, Grimm NB, Howarth RW, Marino R, Martinelli L, Rastetter EB, et al** (2002) Towards an ecological understanding of biological nitrogen fixation. *Biogeochemistry* **57**: 1–45
- West JB, HilleRisLambers J, Lee TD, Hobbie SE, Reich PB** (2005) Legume species identity and soil nitrogen supply determine symbiotic nitrogen-fixation responses to elevated atmospheric CO₂. *New Phytol* **167**: 523–530
- Whittaker JB** (1999) Impacts and responses at population level of herbivorous insects to elevated CO₂. *Eur J Entomol* **96**: 149–156
- Winkler JB, Herbst M** (2004) Do plants of a semi-natural grassland community benefit from long-term CO₂ enrichment? *Basic Appl Ecol* **5**: 131–143
- Zanetti S, Hartwig UA, Luscher A, Hebeisen T, Frehner M, Fischer BU, Hendrey GR, Blum H, Nosberger J** (1996) Stimulation of symbiotic N₂ fixation in *Trifolium repens* L under elevated atmospheric pCO₂ in a grassland ecosystem. *Plant Physiol* **112**: 575–583
- Zavala JA, Casteel CL, DeLucia EH, Berenbaum MR** (2008) Anthropogenic increase in carbon dioxide compromises plant defense against invasive insects. *Proc Natl Acad Sci USA* **105**: 5129–5133
- Zavala JA, Casteel CL, Nabity PD, Berenbaum MR, DeLucia EH** (2009) Role of cysteine proteinase inhibitors in preference of Japanese beetles (*Popillia japonica*) for soybean (*Glycine max*) leaves of different ages and grown under elevated CO₂. *Oecologia* **161**: 35–41
- Ziska LH, Palowsky R, Reed DR** (2007) A quantitative and qualitative assessment of mung bean (*Vigna mungo* (L.) Wilczek) seed in response to elevated atmospheric carbon dioxide: potential changes in fatty acid composition. *J Sci Food Agric* **87**: 920–923