



REVIEW PAPER

# How can we make plants grow faster? A source–sink perspective on growth rate

Angela C. White<sup>1,\*</sup>, Alistair Rogers<sup>2</sup>, Mark Rees<sup>1</sup> and Colin P. Osborne<sup>1\*</sup>

<sup>1</sup> Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

<sup>2</sup> Biological, Environmental and Climate Sciences Department, Brookhaven National Laboratory, Upton, NY 11973, USA

\* To whom correspondence should be addressed. E-mail: [angela.white@sheffield.ac.uk](mailto:angela.white@sheffield.ac.uk); [c.p.osborne@sheffield.ac.uk](mailto:c.p.osborne@sheffield.ac.uk).

Received 7 August 2015; Revised 21 September 2015; Accepted 22 September 2015

Editor: Christine Raines

## Abstract

**Growth is a major component of fitness in all organisms, an important mediator of competitive interactions in plant communities, and a central determinant of yield in crops. Understanding what limits plant growth is therefore of fundamental importance to plant evolution, ecology, and crop science, but each discipline views the process from a different perspective. This review highlights the importance of source–sink interactions as determinants of growth. The evidence for source- and sink-limitation of growth, and the ways in which regulatory molecular feedback systems act to maintain an appropriate source:sink balance, are first discussed. Evidence clearly shows that future increases in crop productivity depend crucially on a quantitative understanding of the extent to which sources or sinks limit growth, and how this changes during development. To identify bottlenecks limiting growth and yield, a holistic view of growth is required at the whole-plant scale, incorporating mechanistic interactions between physiology, resource allocation, and plant development. Such a holistic perspective on source–sink interactions will allow the development of a more integrated, whole-system level understanding of growth, with benefits across multiple disciplines.**

**Key words:** Carbon, crops, models, nitrogen, plant growth, regulation, sink, source.

## Introduction

Growth rates of plants vary widely: even in constant environmental conditions, relative growth rate can vary six-fold among species (Grime and Hunt, 1975). This is not surprising given the astonishing variety of ecological niches occupied by plants in all the major biomes, where adaptation comes in part from matching growth rate to available resources (Díaz *et al.*, 2004). Growth is controlled by proximate physiological and developmental mechanisms, but ultimately depends upon ecological adaptations and evolutionary history: plants with different growth strategies succeed in different ecosystems, and in different niches within those ecosystems. For example, in the dynamic, diverse rainforest environment, rapidly growing seedlings and lianas will quickly colonize gaps, while slow-growing

epiphytes often stay poised and wait for a gap in the canopy before upregulating their rates of photosynthesis and growth (Hubbell and Foster, 1992). Ecological life history theory points towards a growth-survival trade-off (e.g. Baraloto *et al.*, 2010), which helps to explain species differences in growth rate (Metcalf *et al.*, 2006), and leads to niche partitioning (Wright *et al.*, 2010). Growth rate therefore represents a major axis of ecological variation among species, which correlates with changes in resource availability and risk of mortality, but trades off against defence and storage (Grime, 1977; Herms and Mattson, 1992; Rose *et al.*, 2009; Turnbull *et al.*, 2012).

Proximate causes of growth rate variation include both external and internal factors (Körner, 1991). Externally,

plants are affected by a plethora of abiotic and biotic factors including nutrient and light levels, temperature, competition, and herbivory, all of which influence the supply and demand for essential resources, and plants must ensure that growth rates are attuned accordingly (Bloom *et al.*, 1985; Coley *et al.*, 1985). Internally, plant growth is constrained by molecular, physiological, and developmental processes: metabolic rates determine the capacity to take up and store resources, while allocation during development, rates of cell division and expansion, and developmental transitions from vegetative to reproductive growth all have important effects on resource use and partitioning. These internal processes can all be understood within the framework of source–sink interactions: source activity refers to the rate at which essential external resources are acquired by the plant and made available internally, while sink activity refers to the internal drawdown of these resources. This drawdown encompasses resource sequestration in growth and storage, plus resource losses through respiration or exudation. By necessity, the relationships between sinks and sources are both finely tuned and tightly regulated by feedback and feedforward mechanisms, many of which are now well characterized within tissues at the molecular level (e.g. Smith and Stitt, 2007; Lawlor and Paul, 2014). Since plants are sessile and can only influence external factors to a limited degree, the internal factors are well controlled. As a consequence, it is these internal interactions of source and sink activity that must be responsible for the large intrinsic variation in relative growth rate among species under common environmental conditions.

The general principles governing the diversity in intrinsic growth rate among wild species also underpin the variation in yield potential among crop genotypes. The current need to increase crop productivity for food and fuel, due to a rapidly increasing global population, is urgent and well-documented (FAO *et al.*, 2014). Yield increases in rice and wheat due to breeding and genetic techniques are currently around 1% per year—a trajectory too low to meet future requirements, and this has motivated the development of global consortia for crop improvement (von Caemmerer *et al.*, 2012; Reynolds *et al.*, 2012; Ort *et al.*, 2015). The primary focus for many of these is boosting photosynthetic carbon acquisition (source activity), yet sink activity is also believed to limit grain development in many major crops (Acreche and Slafer, 2009; Peterhansel and Offermann, 2012; Slewinski, 2012). Global efforts to elucidate the responses of crop photosynthesis and yield to future elevated atmospheric CO<sub>2</sub> conditions show that the translation of a large and sustained stimulation of photosynthesis into growth and yield differs markedly between species and often falls short of the expected response (Long *et al.*, 2006a). Achieving future yield targets requires that this translation of improved photosynthesis into yield is made effectively through enhanced sink development. To achieve the 70% increase in crop productivity required by 2050, a greater understanding of the relationships between photosynthesis and growth, and the factors underpinning growth rates, is therefore essential.

This review discusses current understanding of plant growth rates, considering a range of factors from molecular

to ecological, with a particular focus on source–sink interactions. It emphasises the importance of sources and sinks as determinants of growth and as targets for crop improvement. For the first time, this review argues the case for a fully integrated network analysis of physiology, allocation, and development when considering growth across the diversity of wild and crop plants. It highlights source and sink limitation as key areas where understanding could be improved, and suggests that quantitative estimates are required of how sources and sinks limit growth, the extent to which these limitations change at different stages of development within the same species, and their differences in species, which vary in allocation and life history strategies.

## Source and sink definitions

Source tissues are net exporters of an elemental resource required for plant growth, such as carbon or nitrogen, while sink tissues are net importers and are responsible for resource assimilation. Mature leaves are net sources of carbon but sinks for nitrogen, while root tissues are net sources of nitrogen but sinks for carbon. Cells require carbon and nitrogen for growth and development; nitrogen to maintain protein turnover; and carbon for respiration to fuel metabolic processes. Other elements are also vital for growth, such as oxygen obtained from the air, hydrogen from water, and minerals found in soil including the macronutrients potassium and phosphorus, and numerous micronutrients. This review focuses on carbon and nitrogen only, because these elements are commonly limiting for growth, and effectively illustrate the balance between source and sink tissues. Carbon is usually exchanged between sources and sinks as simple sugars, typically sucrose. The equivalent currency of exchange for nitrogen includes both inorganic ions (NO<sub>3</sub><sup>−</sup>) and organic forms (typically amino acids).

Source tissues are generally responsible for the acquisition of resources from the external environment, although the remobilization of stored resources (e.g. to subsidize reproduction or regrowth after disturbance) may also turn a sink into an internal source. A general definition of source strength should therefore consider the export rate of a particular resource from the source tissue. However, C- or N-uptake from the external environment is more commonly and easily measured than internal fluxes of sucrose or inorganic and organic nitrogen. Consequently, the term ‘source strength’ usually refers to the net rate of uptake (mol s<sup>−1</sup>) for a particular resource from the external environment:

$$\text{Source strength} = \text{source size} \times \text{source activity} \quad (1),$$

where source size refers to the total biomass of source tissue (g), and source activity is the specific uptake rate of the resource (mol g<sup>−1</sup> s<sup>−1</sup>; based on Geiger and Shieh, 1993).

Sink tissues are net receivers of resources from source tissues (Doehlert, 1993). While all tissues have some sink activity, leaves are net sinks for nitrogen transported from the root system, and roots are net sinks for sucrose exported from leaves. Sink strength refers to the net rate of uptake (mol s<sup>−1</sup>) for a particular resource by a defined tissue within the plant:

$$\text{Sink strength} = \text{sink size} \times \text{sink activity} \quad (2),$$

where sink size is the total biomass of sink tissue (g), and sink activity refers to the specific uptake rate of the resource ( $\text{mol g}^{-1} \text{s}^{-1}$ ). Sink activity involves the utilization of resources for the synthesis of new tissues, including the synthesis of structural components such as cell walls, or the maintenance and modification of existing tissues, including the synthesis of non-structural components including enzymes, storage and defence compounds. Sink activity also encompasses the expenditure of resources in respiration or root exudation. In practice, therefore, it is usually quantified via the net accumulation rate of a particular resource in a tissue over time, after accounting for the losses from respiration and exudation.

Source tissues thus take up resources from the environment and export them to sinks, which draw down resources within the plant. The parallels with financial transactions are clear in this conceptualization of plant function, and the next section considers the molecular currencies traded between sources and sinks.

## Source and sink tissues

### Carbon

Mature leaves are net sources of carbon. Carbon dioxide is fixed to generate triose phosphate in photosynthesis, which is then converted to starch for diurnal storage in the chloroplast (Smith and Stitt, 2007; Gibon *et al.*, 2009; Stitt and Zeeman, 2012; Pilkington *et al.*, 2015), or to sucrose for export from the leaf or storage in the vacuole.

Net carbon sink tissues include roots, tubers, reproductive structures, and young leaves. Sucrose may itself be stored directly, or it may first be converted to storage polymers. These polymers are typically starch or fructans, depending on the species; some plants store additional compounds such as raffinoses (Atkinson *et al.*, 2012). Starch is stored in the amyloplasts and chloroplasts of many higher plants (Müller-Röber *et al.*, 1992); amyloplasts are found in seeds, shoot storage tissues, and roots, while chloroplasts are found in leaves and stems (and are the only repository for starch within leaves). Starch is also the primary carbohydrate in the grains of many crops, including wheat, rice, and maize, and in the tubers and storage roots of vegetables (Pollock and Cairns, 1991; Zeeman *et al.*, 2010). Carbon storage in the stems of many temperate grasses consists primarily of fructans (Pollock and Cairns, 1991; Scofield *et al.*, 2009), water-soluble fructose polymers which confer some resistance to low temperatures (Sandve *et al.*, 2011). Typically, fructans and sucrose are stored together in the stem, as in wheat, barley, and oat (Slewinski, 2012). Significant stem storage of starch is rare in cereals—rice being a notable exception, storing sucrose in leaves and starch in stems (Murchie *et al.* 2009), and being unable to synthesize fructans naturally (Kawakami *et al.*, 2008). Stem storage of carbohydrates is an important buffering system for recovery after grazing and for supplying photosynthate to cereal ears during grain-filling, especially during drought

(Schnyder, 1993; Ruuska *et al.*, 2006; Slewinski, 2012), and is thus a relatively labile sink. For growth, a major use of photosynthate is the synthesis of cell wall polysaccharides such as cellulose and hemicellulose, in all parts of the plant. Indeed, almost half of plant cell wall biomass is composed of carbon (Körner, 2012).

In addition to the assimilation of resources in sink tissues, the utilization of resources in respiration and exudation constitute a further sink, since these processes also contribute to resource drawdown. Maintenance respiration can represent a significant carbon cost to the plant (Penning de Vries, 1975); for example, respiration constitutes 70% of the carbon sink in *Pinus halepensis* (Klein and Hoch, 2015). Carbon and nitrogen are released through root exudation of a variety of compounds including organic acids, sugars, polysaccharides, ectoenzymes such as acid phosphatase, and sloughed-off cells and tissues (Marschner, 1995). Exuded metabolites have many functions (Badri and Vivanco, 2009) such as modifying the rhizosphere to provide a desirable environment for beneficial microorganisms and providing signals to aid recruitment of arbuscular mycorrhizal fungi, contributing to immunity (Cameron *et al.*, 2013). These processes may be substantial—one meta-analysis of annuals found that 30–60% of net photosynthetic carbon is allocated to roots, of which 40–90% is lost in respiration and exudation (Lynch and Whipps, 1990).

### Nitrogen

In contrast to carbon, roots are net sources for nitrogen, while shoot tissues are net nitrogen sinks until senescence when their nitrogen is remobilized (Aerts and Chapin, 2000). Annuals remobilize nitrogen for reproduction while perennials may remobilize nitrogen for reproduction or for growth and storage in subsequent years. Inorganic nitrogen is taken up by roots as nitrate ( $\text{NO}_3^-$ ) or ammonium ( $\text{NH}_4^+$ ), and may be utilized in growth or exported from the root. Assimilation of nitrogen into amino acids takes place in both roots and shoots, although the relative proportions depend on the species and are still debated (Nunes-Nesi *et al.*, 2010). Approximately 80% of wild plant species benefit from mycorrhizal associations in which specialised fungi aid the uptake of phosphorus and sometimes organic nitrogen (Read, 1991). Organic nitrogen may also be taken up from the soil in the form of free amino acids. Nitrogen is exported from roots as nitrate (transported in the xylem), amino acids or amides (both transported in the phloem).

Root and leaf nitrogen concentrations are positively correlated, but a global survey of wild grassland species found that leaf nitrogen concentration is more than double that of roots in the same species (Craine *et al.*, 2005). The main use of nitrogen for growth is in proteins and there is a particularly high demand in leaves, where the complex, enzyme-rich photosynthetic machinery is assembled and maintained. Photosynthetic proteins encompass the majority of leaf nitrogen—for example, Rubisco (EC 4.1.1.39) typically accounts for between 10% and 30% of leaf nitrogen content but can account for up to 50% of leaf nitrogen content (Ellis, 1979; Sage *et al.*, 1987; Evans, 1989). Through Rubisco, carbon

source activity is directly connected with leaf nitrogen sink activity, providing one way in which source and sink activity are intrinsically coordinated. Nitrogen partitioning into photosynthetic proteins is a flexible trait, varying between species and as resource availability changes (Evans, 1989).

Nitrogen may be stored as nitrate in the vacuole, or as proteins (Millard, 1988). Vegetative storage proteins (VSPs) may comprise up to 50% of soluble protein in vegetative tissues (Liu *et al.*, 2005). Protein storage occurs primarily in seeds, although some legumes, tuber-formers, and deciduous trees species have additional storage proteins (Shewry, 1995). VSPs have been well-studied in soybean, and the nitrogen sink-to-source transition occurring in leaves during ontogeny is correlated with a decrease in VSP gene expression in this species (reviewed by Staswick, 1990). In both potato and soybean, removal of nitrogen sink tissues upregulates nitrogen storage in other parts of the plant, indicating a buffering role for VSPs in maintaining source:sink balance (Staswick, 1990). In contrast, grasses such as wheat and rice (usually grown as annuals) are less reliant on protein stores beyond those in the seed, yet can still accumulate nitrogen when conditions are favourable. For example, excess nitrogen in wheat accumulates in the lamina of upper leaves or the true stem of the peduncle, and just as stem carbohydrate reserves are important for grain-filling in grasses, this stored nitrogen is thought to provide a nitrogen buffer for use during grain-filling (Pask *et al.*, 2012). Non-leaf nitrogen stores, such as the culm in grasses, may also play a role in plant recovery after grazing.

## Co-limitation and optimization

A plant that has optimized its source:sink ratio can grow using balanced allocation of source and sink sizes and activities (Equations 1–2), facilitated by molecular feedbacks. However, at any point in time, most plants are not fully optimized, meaning that the creation of either more source or sink tissue could increase growth: in these cases, the potential source or sink strength (Equations 1–2) has not been realized (Patrick, 1993). The extent to which these potentials are met may be investigated using environmental or genetic manipulations, and are discussed later in this review.

Resource uptake changes over time due to fluctuations in the external environment, such that the total supply of a particular resource over the lifetime of the plant cannot be predicted in advance. The plant reacts to these fluctuations in resource availability by modifying its investment in resource acquisition and consumption (Freschet *et al.*, 2015). At the most general level, a suitable balance between leaf and root tissues is critical for balancing the acquisition of carbon and mineral nutrients. Allocation to shoot and root is adjusted depending on available resources so that, for example, the allocation of resources to root growth is increased in low nitrogen soil conditions. Co-limitation by carbon and nitrogen has been demonstrated experimentally and optimization of these resources has been considered in models (Woodrow, 1994; Iwasa, 2000; Guilbaud *et al.*, 2015) yet, due to environmental

and developmental constraints, plants do not always achieve perfect co-limitation *in vivo*.

Greater insights into this balancing of sources and sinks at the whole-plant scale can be gained by analogy with metabolic systems within cells or tissues. In plant metabolic networks, control of the overall flux is typically shared between several enzyme steps, although many elements in the system exert only a limited effect (Fell and Thomas, 1995; e.g. Raines, 2003; Araújo *et al.*, 2012). If the dynamic internal system of resource fluxes among source and sink tissues is analogous to such a metabolic system, then overall control of the flux of materials into growth is also likely to be shared among multiple elements. This flux control analogy generates two predictions.

The first prediction is that multiple elements in the system share control of the growth rate, and growth will increase if their sizes or activities are raised together. In contrast, most elements exert limited control, and are present in excess. The most resource-efficient solution for the developing plant is therefore to tune down investment in those components with little influence, and increase allocation to the elements exerting a high degree of control. Such regulation must be a dynamic process that balances fluctuations in external resource availability with ontogenic changes in the demand for resources. Analogous examples from metabolism show how such reallocation among elements in the system can optimize enzyme activities to increase fluxes (Woodrow, 1994; Zhu *et al.*, 2007). However, in a whole-plant system, this optimization process must operate within the context of life history strategies of investment in growth versus defence or storage (the growth–survival trade-off).

The second prediction generated by the flux control analogy is that development of new source and sink organs during ontogeny shifts the overall control of growth to different elements in the system. This effect is expected because changes in the number, size, and activity of plant organs during development alters the internal capacity of a plant to acquire and consume resources, and is well supported by experimental evidence. For example, some plants transition from sink to source limitation during the shift from vegetative to reproductive growth (examples within Arp, 1991; Marschner, 1995; Rogers and Ainsworth, 2006). Equivalent effects arise when plants are exposed to external environmental conditions which change resource acquisition rates, or if the numbers or activities of source or sink organs are manipulated experimentally. The evidence for such effects is considered in the next section of this review.

In combination, these external and internal factors mean that the acquisition and consumption of resources must be balanced over time by a combination of coarse and fine internal regulatory controls. This control, in turn, operates within a general life history strategy of investment in growth versus storage or defence, which means that the growth rate is not necessarily maximized under particular internal and external constraints.

The situation for crop plants is simpler, since breeders aim to maximize lifetime growth and reproductive allocation within monospecific communities (Denison, 2012). Current

views on source–sink relations in crop plants point towards a co-limitation of growth by sources and sinks during grain-filling (Álvarez *et al.*, 2008; Acreche and Slafer, 2009; Peterhansel and Offermann, 2012; Slewinski, 2012) yet growth could be further optimized. One line of evidence for the lack of optimization of source and sink to maximize growth comes from experiments where plants are grown at elevated CO<sub>2</sub> (discussed later, in Table 1). Such experiments aim to predict the responses of plants to future climatic conditions, and increase the carbon source activity of plants in a non-invasive manner. The increase in photosynthesis under elevated CO<sub>2</sub> demonstrates that source activity typically limits growth under ambient CO<sub>2</sub> levels. However, the increases in photosynthesis and yield seen when plants are grown under elevated CO<sub>2</sub> do not match the magnitude of those predicted from theoretical modelling and extrapolation of chamber experiments (Long *et al.*, 2006a; Ainsworth *et al.*, 2008; Leakey *et al.*, 2009). These results suggest a degree of sink limitation of growth, which could be due to nitrogen limitation. Responses to CO<sub>2</sub> do vary between species (Poorter, 1993) and some plants are able to upregulate source and sink in concert. For example, high CO<sub>2</sub> can stimulate nitrate uptake to balance source and sink capacity (Stitt and Krapp, 1999). When external nitrate levels are low, elevated CO<sub>2</sub> levels cause an increase in both the rate of nitrate uptake and the activity of a high affinity nitrate transport system in wheat roots (Lekshmy *et al.*, 2009), representing an upregulation of nitrogen source strength through increased activity [Equation (1)].

In order to improve crop yields, a greater, more integrated understanding of how plant growth rates are limited by sinks and sources for carbon and nitrogen, and the shifts in limitation that occur during the lifetime of a plant, is required. Only by grounding modelling and experimental work in mechanistic knowledge of source:sink relationships will plant growth be effectively understood—and potentially manipulated—at every stage of development in order to maximize yield.

## Sources and sinks affect growth and yield

Evidence that growth may be controlled by both source and sink strengths comes from manipulation experiments and studies of natural variation among species.

### *Manipulation experiments*

Manipulating the source:sink balance shows that source and sink strengths often operate below their full potential, due to the limitations imposed by environmental and developmental changes discussed above. Historically, such manipulations involved physically manipulating the plant or its environment: for example, source activity may be altered by elevated CO<sub>2</sub>, defoliation, or shading, while sink activity may be altered by sink removal or sink chilling. However, modern genetic approaches may now be used to alter source and sink activity with greater elegance. Table 1 outlines a range of source:sink manipulations and summarizes their results. Broadly speaking, increasing either source or sink may increase growth,

suggesting that both can limit growth to a certain extent. Sources and sinks regulate each other by molecular feedback mechanisms (discussed later), and evidence for these is seen at the whole-plant scale when manipulation of the source affects sink activity, and *vice versa*.

Elevated CO<sub>2</sub> increases the potential carbon source activity of the plant by stimulating photosynthesis, and this typically translates into faster growth (Table 1; e.g. McConnaughay *et al.*, 1993; Christ and Körner, 1995; Masle, 2000; see also Taylor *et al.*, 1994; Ranasinghe and Taylor, 1996; Long *et al.*, 2006b; Ainsworth and Rogers, 2007; Leakey *et al.*, 2009), also affecting cell patterning, cell expansion, and plant architecture (Kinsman *et al.*, 1997; Pritchard *et al.*, 1999; Masle, 2000). Growing plants in large pots increases the potential carbon sink activity, due in part to increased nitrogen availability, generally leading to increased growth (Table 1; McConnaughay *et al.*, 1993; Poorter *et al.*, 2012), and experiments comparing species or cultivars with different sink sizes reveal that growth is faster when sinks are larger (Table 1; Reekie *et al.*, 1998; Aranjuelo *et al.*, 2013).

Reduction of source leaf area by defoliation usually leads to an increase in photosynthesis in the remaining leaves, to maintain source activity within the plant and support the sinks (Table 1; von Caemmerer and Farquhar, 1984; Eyles *et al.*, 2013). This could indicate sink limitation of growth because leaves are not carrying out their maximal potential rates of photosynthesis under normal conditions. In contrast, decreasing sink capacity—for example, by inhibiting sucrose export from leaves to reduce the apparent sink demand—leads to an inhibition of photosynthesis (Table 1; Ainsworth and Bush, 2011) mediated by an increase in leaf carbohydrates (Sheen, 1990).

Combining experimental treatments that affect both source and sink provides evidence that sources and sinks work together and feed back on each other. Photosynthetic acclimation at elevated CO<sub>2</sub> concentration is a decrease in photosynthetic capacity that reduces the magnitude of the CO<sub>2</sub>-induced stimulation in photosynthetic rate at elevated CO<sub>2</sub>. Acclimation acts to reduce the ratio of source:sink activity and thus adjust source:sink balance towards equilibrium. Combining defoliation and elevated CO<sub>2</sub> treatments (which decrease and increase the source, respectively) shows that photosynthetic acclimation under elevated CO<sub>2</sub> is alleviated by defoliation, supporting the hypothesis that it is sink-mediated (Table 1; e.g. Bryant *et al.*, 1998; Rogers *et al.*, 1998; Ainsworth *et al.*, 2003). The alleviation of acclimation occurs because defoliation opposes the effect of elevated CO<sub>2</sub> by decreasing the source:sink ratio, and higher levels of photosynthesis can thus be maintained in the remaining leaves. The opposite effect is seen when the source is increased but the sink is reduced. For example, when physical removal / restriction of sinks or genetic manipulation to reduce sink size is combined with elevated CO<sub>2</sub>, leading to an increase in the source:sink balance, source activity is decreased in order to return towards equilibrium (Table 1; Arp, 1991; Ainsworth *et al.*, 2004). Combining low nitrogen or low temperature—which restrict sink development—with elevated CO<sub>2</sub> has a similar effect (Table 1; Arp, 1991). In contrast, increasing

**Table 1.** Experimental manipulations of the carbon source:sink balance, illustrating that: (a) both sources and sinks affect plant growth; (b) sources and sinks regulate each other by feedback mechanisms; (c) source and sink strength can be altered by the plant, to alleviate perturbations of the source:sink balance

Species	Manipulation	Effect	Key result	Reference
SOURCE MANIPULATIONS				
<i>Eucalyptus globulus</i>	Defoliation +	Reduces source	Defoliation increases photosynthesis in other leaves; source:sink biomass ratio is main driver of this change	(Eyles et al., 2013)
Three chalk grassland species	Debudding Defoliation +	Reduces sink Reduces source	In two species, photosynthetic acclimation to elevated CO <sub>2</sub> was alleviated by defoliation, which restores the source:sink balance	(Bryant et al., 1998)
	Elevated CO <sub>2</sub>	Increases source		
<i>Lolium perenne</i>	Canopy-cutting +	Reduces source	Photosynthetic acclimation to elevated CO <sub>2</sub> was alleviated by cutting the canopy, which restores the source:sink balance	(Rogers et al., 1998)
	Elevated CO <sub>2</sub>	Increases source		
<i>Phaseolus vulgaris</i>	Defoliation / Reduced light +	Reduces source	At ambient and elevated CO <sub>2</sub> : defoliation increases photosynthetic rate in other leaves; reduced light decreases photosynthetic rate	(von Caemmerer & Farquhar, 1984)
	Elevated CO <sub>2</sub>	Increases source		
<i>Lolium perenne</i>	Elevated CO <sub>2</sub> +	Increases source	Photosynthetic rate decreased in low nitrogen, but this effect was reduced when the source:sink balance was restored by	(Ainsworth et al., 2003)
	Canopy-cutting +	Reduces source	canopy-cutting	
	Low nitrogen	Reduces sink		
<i>Dactylis glomerata</i>	Elevated CO <sub>2</sub>	Increases source	Shortening of cell cycle in shoot and root meristems	(Kinsman et al., 1997)
<i>Triticum aestivum</i>	Elevated CO <sub>2</sub>	Increases source	Cell division and expansion affected	(Masle, 2000)
SINK MANIPULATIONS				
Various species	Elevated CO <sub>2</sub> +	Increases source	Reducing sink capacity increases acclimation of source activity	(Arp, 1991)
	Removal of sinks / Low nitrogen / Low temperature	All reduce sink		
<i>Arabidopsis thaliana</i>	Low temperature +	Reduces sink	Altered signalling pathway reduced plant capacity to recover from sink limitation	(Nunes et al., 2013)
	Genetic manipulation of T6P/ SnRK1 signalling pathway	Affects integration of sucrose levels and growth		
Various species	Inhibition of sucrose export from source leaves	Reduces apparent sink demand	Inhibition of photosynthesis	(Ainsworth and Bush, 2011)
<i>Glycine max</i>	Elevated CO <sub>2</sub> +	Increases source	Reduced sink capacity and decreased photosynthesis, due to increase in source:sink balance	(Ainsworth et al., 2004)
	Genetic modification to make a determinate line of a cultivar normally showing indeterminate growth	Reduces sink		
<i>Solanum tuberosum</i>	Transgenic reduction of ADP-glucose pyrophosphorylase	Reduces sink capacity by reducing starch synthesis	Tuber sinks adapted by increasing sucrose content	(Müller-Röber et al., 1992)
<i>Solanum tuberosum</i>	Transgenic reduction of ADP-glucose pyrophosphorylase +	Reduces sink capacity by reducing starch synthesis	Plants avoided yield reductions by synthesizing fructan instead	(Zuther et al., 2011)
	Transgenic expression of fructan biosynthesis enzymes	Increases sink		
<i>Triticum aestivum</i>	Transgenic modification to increase sucrose uptake in developing grains	Increases sink	Storage protein synthesis increased	(Weichert et al., 2010)

Table 1. Continued

Species	Manipulation	Effect	Key result	Reference
<i>Triticum aestivum</i>	Elevated CO <sub>2</sub> +	Increases source	Acclimation of photosynthesis did not occur when nitrogen was added in this way	(Farage <i>et al.</i> , 1998)
	Addition of nitrogen in proportion to growth	Increases sink		
<i>Abutilon theophrasti</i> and <i>Setaria faberii</i>	Elevated CO <sub>2</sub> +	Increases source	Increase in growth and yield in response to elevated CO <sub>2</sub> was higher when sink capacity was also increased	(McConnaughay <i>et al.</i> , 1993)
	Large size / High nutrients	Both increase sink		
<i>Triticum aestivum</i>	Elevated CO <sub>2</sub> +	Increases source	Increase in photosynthesis and growth was dependent on high sink strength: only seen in cultivar with high harvest index	(Aranjuelo <i>et al.</i> , 2013)
	Cultivars with high and low harvest index	Different sink sizes		
<i>Brassica</i> spp.	Elevated CO <sub>2</sub> +	Increases source	Long-term growth increases were dependent (to an extent) on species-specific sink size	(Reekie <i>et al.</i> , 1998)
	Species had different sink sizes	Different sink sizes		

'+' denotes treatments applied in combination; '/' denotes alternative treatments.

carbon sink capacity under elevated CO<sub>2</sub>, by using high-yielding cultivars or adding nitrogen, facilitates increased photosynthesis (Table 1; Farage *et al.*, 1998; Aranjuelo *et al.*, 2013).

#### Differences among species

In some species, developmental plasticity allows for greater flexibility when the source:sink balance is perturbed. Potato and citrus may easily increase their sink size, so tend to suffer less from feedback inhibition of photosynthesis (Paul and Foyer, 2001), and nitrogen-fixing legumes are easily able to increase their sink size in response to elevated CO<sub>2</sub> (Rogers *et al.*, 2009).

The physical mechanism of carbon export is important for the coordination of source and sink. Growth determinacy in soybean prevents an increase in photosynthesis at high CO<sub>2</sub>, while poplar trees have high photosynthate export and maintain elevated photosynthesis at high CO<sub>2</sub> (Table 1; Ainsworth *et al.*, 2004; Ainsworth and Bush, 2011). Species which are symplastic loaders (many trees and shrubs) transport sucrose from source tissues into the phloem through developmentally fixed plasmodesmata, whereas apoplastic loaders (many herbaceous species) use developmentally plastic membrane transporters (Ainsworth and Bush, 2011). Therefore, at high CO<sub>2</sub>, symplastic loaders cannot upregulate photosynthate export to the same extent as apoplastic loaders. As a result they tend to accumulate more non-structural carbohydrates in their leaves (Körner *et al.*, 1995) and can show a smaller increase in photosynthesis under elevated CO<sub>2</sub>. However, despite their symplastic loading strategy, trees are generally well able to maintain photosynthetic stimulation under elevated CO<sub>2</sub> (Ainsworth and Rogers, 2007) although some species are capable of both symplastic and apoplastic loading and many species have not yet been characterized.

Taken together, this evidence clearly demonstrates that sources and sinks can both limit growth, and that feedbacks enable a degree of compensation. Species with greater plasticity can be more flexible in their response to manipulations of source and sink.


## Regulation of sinks and sources

Regulation of source:sink balance is essential for enabling plants to maintain a growth rate appropriate for a given availability of resources. Storage allows the assimilation of more resources than are needed in growth, to create a reserve for future development in a fluctuating environment or recovery from disturbances such as herbivory. However, carbon assimilation must be appropriate for the available sink strength, in order to create a sufficiently large store that is still within the limits imposed by sink potential—thus sinks must feed back on sources to regulate their activity. Similarly, source activity must influence sink strength so that appropriate sinks may develop and plants can fully realize their growth potential for a given resource availability. Furthermore, the high metabolic costs of carbon and nitrogen assimilation mean that regulation of source and sink is vital to avoid wasting energy.

A complex molecular network including carbon- and nitrogen-derived signals and phytohormones has evolved to integrate the uptake, assimilation, and allocation of resources (Nunes-Nesi *et al.*, 2010). Many mechanisms of these molecular interactions are now well established although the puzzle remains incomplete at the whole-plant scale. Figure 1 illustrates key feedforward and feedback mechanisms regulating the source:sink relationship. Carbon- and nitrogen-derived feedforward and feedback signals act on sources and sinks of both carbon and nitrogen. This allows sources and sinks to modify their own activity, and also to regulate that of other tissues, creating molecular signalling links between source and sink.

#### Carbon feedbacks

Leaf carbohydrates feed into a complex network, affecting transcription, translation, and post-translational processes in order to balance carbon supply and demand (reviewed in Fig. 1). For example, a high carbon status upregulates nitrogen source and sink activity (Fig. 1; arrows 7 and 9) and carbon sink activity (arrows 5a and 5c), while downregulating photosynthesis

**KEY:**

1. If sink strength for carbon is high, **nitrogen-derived signals** increase carbon source activity. **Cytokinins** travel from root to shoot (10) and upregulate photosynthetic genes when the root nitrogen concentration – and thus the capacity for carbon usage – is high (Paul and Foyer, 2001). **Nitrogen-derived signals** also modulate feedback of **sugars** on photosynthesis (2) (Stitt and Krapp, 1999).
2. When **sugars** accumulate in the leaf they repress the expression of photosynthetic genes (reviewed by Smith & Stitt, 2007); the glucose sensor **hexokinase** leads to stomatal closure when sucrose production exceeds drawdown by phloem loading and thus reduces photosynthesis (Kelly et al., 2013); **carbohydrate-accumulation**, **redox signals**, and **phosphate recycling** enable sink regulation of photosynthesis in the short term (Paul and Foyer, 2001).
3. **Sugars** in the phloem regulate export of leaf carbohydrates (Ainsworth and Bush, 2011). Inhibition of sucrose loading can increase the concentration of sucrose in the leaf, leading to a reduction in source activity (2). For example, a proton-sucrose symporter in sugar beet is regulated by sucrose, allowing high sucrose levels in the phloem to influence export from the leaf by negative feedback (Vaughn et al., 2002).
4. **Cytokinins** increase cell division and sink strength, increasing growth when nitrogen is plentiful (Kuiper, 1993; Ghanem et al., 2011; Thomas, 2013).
- 5a. The glucose sensor **hexokinase**, the primary metabolite **trehalose-6-phosphate (T6P)** and the **TOR protein kinase signalling pathway** provide three major ways in which sugar levels upregulate plant growth (Smeekens et al., 2010). Sucrose levels upregulate **T6P** and **TOR**; **TOR** increases growth and **T6P** appears to be an important integrator of sucrose and growth although its role is still debated (Xiong et al., 2013; Nunes et al., 2013; Lastdrager et al., 2014; Lunn et al., 2014). This leads to a positive feedback between source activity and sink activity.
- 5b. The **protein kinase SnRK1** and the **C/S1 bZIP transcription factor network** inhibit growth (reviewed by Lastdrager et al., 2014; Smeekens et al., 2010) and are regulated by metabolite status, allowing source to repress sink by reducing growth when carbon resources are scarce.
- 5c. **Sucrose** and **T6P** repress the action of **bZIP transcription factors** and **SnRK1** respectively, lifting repression of growth (5b) when photosynthate is plentiful (Lastdrager et al., 2014).
6. **Nitrate** and **cytokinins** increase transcription of nitrate reductase (NR), a key enzyme in nitrogen assimilation (Stitt and Krapp, 1999). This allows plants to exploit high soil nitrogen. NR transcription is induced by **nitrate** and repressed by downstream metabolites such as **glutamine** (Klein et al., 2000).
7. **Sugars** are thought to positively regulate NR in the root (Stitt and Krapp, 1999).
8. **Nitrate** and **cytokinins** upregulate NR, and thus nitrogen assimilation, in the shoot as in the root (6).
9. **Sugars** regulate shoot NR at a transcriptional and post-translational level; low sugar levels can override control of NR by nitrate and repress NR transcription (Stitt and Krapp, 1999; Klein et al., 2000; Kaiser et al., 2002; Reda, 2015).
10. **Nitrogen** increases **cytokinin** levels and these are exported to the shoot. When plant nitrogen levels are high, **nitrogen-derived signals** have a range of effects at both local (e.g. increasing lateral root growth) and whole-plant (e.g. increasing shoot:root allocation) levels (reviewed by Stitt & Krapp, 1999).

**Fig. 1.** A range of feedback mechanisms fine-tunes the source:sink balance and therefore plant growth. Signals derived from both carbon (green) and nitrogen (blue) regulate source–sink relationships. Feedbacks operate at the tissue level (arrows 1–4 and 6–9) and at the whole-plant level (arrows 5 and 10). Narrow grey arrows represent net movement of carbon and nitrogen from source (lighter) to sink (darker) tissues within the plant.

(arrows 2 and 3). In contrast, a low carbohydrate content in the leaf leads to the repression of carbon sink activity (arrow 5b). The presence of such a regulatory feedback loop in the leaf has long been investigated: in 1868, Bousingault first proposed that assimilate accumulation could decrease photosynthesis by feedback (Neales and Incoll, 1968), yet the precise mechanism for sucrose signalling remains unknown (Reda, 2015).

The partitioning of carbon into starch and sucrose is an important point of carbon source–sink regulation (directly affecting arrows 2 and 5 in Fig. 1) and is controlled by several factors. For example, trehalose-6-phosphate is believed to influence starch synthesis by redox-regulation of AGPase, a key enzyme in starch synthesis, while the degradation of starch is regulated by a variety of enzymes, by the circadian clock, and possibly by starch-derived signals or even the level of starch itself (Smith and Stitt, 2007). It is important to note that most research into sugar and starch regulation has been carried out in *Arabidopsis* and it is therefore necessary to expand current knowledge of regulatory mechanisms in crop plants, which may not share the same mechanisms. For example, the starch degradation pathway in the endosperm of cereal grains differs from that in *Arabidopsis* leaves (Smith, 2012), while mutation of PGM, an enzyme important in starch synthesis and essential for normal growth in *Arabidopsis*, does not affect all

species equally, suggesting the use of different metabolic pathways or storage compounds (Stitt and Zeeman, 2012).

### Nitrogen feedbacks

Just as carbon availability impacts both on carbon and nitrogen source and sink activities, nitrogen availability regulates the uptake and storage of carbon (reviewed in Fig. 1). A high nitrogen status increases the rate of carbon acquisition in photosynthesis and also upregulates carbon sinks (Fig. 1; arrows 1 and 4). Nitrogen also increases the assimilation of nitrate by the enzyme nitrate reductase, to upregulate nitrogen source and sink activity (Fig. 1; arrows 6 and 8), and increases shoot:root allocation, enabling the plant to acquire more carbon and make use of the available nitrogen (arrow 10). Furthermore, nitrate increases root cytokinin production and export (Fig. 1, arrow 10), important for meristem generation and function in shoot and root (Su et al., 2011).

### Crosstalk

Tight control of the source–sink relationship is facilitated by points of crosstalk between carbon- and nitrogen-signalling



pathways. This enables plants to maintain a degree of co-limitation for sources and sinks, and carbon and nitrogen. Starch synthesis is regulated by nitrate as well as by sugar: nitrate downregulates transcription of the gene encoding the regulatory subunit of AGPase, an enzyme involved in starch synthesis. This negative regulation by nitrate lowers starch accumulation and allows more leaf sugar to be exported for growth when nitrate levels are high (discussed by [Stitt and Krapp, 1999](#)). Leaf sugars are involved in the transcription and post-translational regulation of nitrate reductase ([Fig. 1](#), arrow 9), enabling plants to coordinate carbon and nitrogen supply ([Stitt and Krapp, 1999](#); [Kaiser et al., 2002](#)): sugars increase the level of nitrate reductase ([Reda, 2015](#)) while low sugar levels repress its transcription ([Klein et al., 2000](#)).

Coordination is enhanced still further by crosstalk between sugars and phytohormones [for recent review, see [Lastdrager et al. \(2014\)](#)]. This contributes to developmental processes such as meristem activity (which is generally upregulated by cytokinins, e.g. [Fig. 1](#), arrow 4) and lends an added layer of complexity to growth regulation ([Eveland and Jackson, 2012](#)). For example, sugars interact with abscisic acid ([Teng et al., 2008](#)) and with auxin ([Stokes et al., 2013](#)). Sugars can also act directly on development, independently of phytohormones, and are believed to be important for regulating meristem activities ([Eveland and Jackson, 2012](#)). Furthermore, sugar levels influence the transcription of thousands of genes; sugars and the circadian clock regulate each other; and sugars induce phytochrome-interacting factors, which regulate growth ([Lastdrager et al., 2014](#)).

In summary, molecular feedbacks including carbon- and nitrogen-derived signals regulate sources and sinks for carbon and nitrogen. Crosstalk exists both between these signals and with growth regulators. With such elaborate molecular mechanisms in place—and given sufficient resources—increasing the sink activity of a plant might be expected to increase its source activity, and vice versa. However, as discussed above, experimental manipulations of sink strength and of source strength reveal that growth cannot always be altered as expected (e.g. [Long et al., 2006a](#)). It has thus become important to increase knowledge of the potential strengths of source and sink, the limits to their physiological interactions, and to better incorporate known molecular mechanisms of the source–sink relationship into models of whole-plant growth. Moreover, in order to effectively increase crop yield, it may be necessary to manipulate the molecular feedback mechanisms between source and sink, in addition to manipulating the strengths of source and sink themselves. A source–sink-based perspective on growth is therefore an essential cross-disciplinary tool for understanding and increasing the growth and yield of crops.

## Alternative perspectives on growth

Different disciplines have alternative perspectives of plant growth. Advancing the mechanistic understanding of growth that is necessary to realize improvements in crop growth will require a unification of these disciplinary perspectives.

Here, a parsimonious model of plant growth which unites these different perspectives is presented. An extremely simplified system is used for illustration. Various factors have been omitted for simplicity, clarity, and ease of unification. These are both intrinsic (additional resources and tissue types within the plant, and feedbacks between internal processes) and extrinsic (environmental limitations on physiological and developmental processes), since plant growth and development are the product of genetic and environmental processes (e.g. [Prusinkiewicz et al., 2009](#); [Pantin et al., 2012](#)). Rather than provide comprehensive models of growth, this section highlights key processes of interest for each of the three perspectives on growth, and uses equations to demonstrate the focus of each. In each perspective, the processes of interest depend on source and sink activities and tissues, and the equations are finally united to form a basic holistic model of plant growth which is underpinned at every level by source–sink interactions.

Growth may be conceptualized in a number of different ways, and may be viewed through different lenses depending on the perspective adopted. Three classic perspectives on growth are based on: the physiology of resource acquisition and loss; the internal allocation of resources to source and sink organs; and the morphogenetic development of source and sink tissues. Crucially, these three alternative perspectives, adopted by communities of scientists from different disciplines are all readily conceptualized within the context of source–sink interactions.

Here, equations have been used to illustrate each definition of growth, by considering a highly simplified system in which a single resource (carbon) is acquired by a source tissue (leaves) and used by sinks (in both leaves and roots). This system enables the limitations on growth to be formally defined in a readily interpreted form, yet still allows growth to be viewed through the three alternative lenses presented. Each of the three perspectives presented is, by mathematical definition, true. However, each is based implicitly upon an alternative hypothesis about the critical intrinsic controls on growth.

At its most fundamental level, growth may be defined as an increase in plant mass over time. For simplicity, growth is considered equivalent to net organic carbon gain, and the acquisition of other resources is ignored. The dry weight of organic carbon in the plant is  $W_p$ , referred to in this section as mass, and absolute growth rate (AGR) is thus net carbon gain over time, in  $\text{g d}^{-1}$ :

$$\frac{dW_p}{dt} = \text{Absolute growth rate} \quad (3).$$

Growth may now be defined in various ways according to the perspective adopted, but the central definition [Equation (3)] is retained. The different approaches to explaining growth focus attention on different primary limitations.

### Physiology

The first approach is physiological: a flux balance of organic carbon for the plant based on the loss and acquisition of this

essential resource to and from the atmosphere via the processes of respiration and photosynthesis (Lambers *et al.*, 1989; Poorter and van der Werf, 1998).

This carbon-based balance viewpoint on growth is adopted widely in crop production models and in Ecosystem and Earth System Models (EESMs), which simulate the physical properties and carbon exchange of the vegetated land surface (e.g. Knorr, 2000; Sitch *et al.*, 2003; Lu and Ji, 2006; Zaehle and Friend, 2010), and are ultimately used to project future global change (Friedlingstein *et al.*, 2014; IPCC, 2014). This flux balance approach expresses the AGR as the difference between photosynthesis and respiration:

$$\frac{dW_P}{dt} = A \times W_L - R \times W_P \quad (4a),$$

where  $A$  is gross photosynthetic carbon uptake in  $\text{g C d}^{-1} \text{g}^{-1}$  leaf mass,  $W_L$  is total leaf mass (g), and  $R$  is respiratory carbon loss in  $\text{g C d}^{-1} \text{g}^{-1}$  plant mass. Note that not all of the inorganic carbon captured by photosynthesis is converted to biomass, and so  $R$  includes the metabolic costs of biosynthesis, translocation, exudation, and the uptake and assimilation of nitrogen needed for growth ('growth respiration'), as well as those associated with maintaining existing tissues ('maintenance respiration') (reviewed by Amthor, 2000).

Respiration may be partitioned between photosynthetic and non-photosynthetic tissues:

$$\frac{dW_P}{dt} = (A \times W_L) - (R_L \times W_L) - (R_R \times W_R) \quad (4b),$$

where the subscripts  $L$  and  $R$  denote leaf and root tissues, respectively. A simple case is considered here, but this approach may be easily extended to include other sink tissues such as storage organs, stems, and reproductive tissues.

This basic model views growth as the net accumulation of organic carbon. However, the approach is limited because, while respiration is one component of sink activity, the sink activities of growth and storage are not explicitly considered, and accounting for sink limitation requires modifications to the model (Fatichi *et al.*, 2013). Furthermore, recent authors have argued that  $A$  and  $R$  do not control plant growth rate. Instead, it is argued that growth is controlled by the supply of mineral nutrients and water, and the plant regulates  $A$  and  $R$  to meet its growth requirements (Körner, 2012; Fatichi *et al.*, 2013; Körner, 2013). Without accounting for sink activities and their feedbacks on photosynthesis, the approach illustrated by Equations (4a) and (4b) cannot provide a complete description of the processes controlling growth.

### Allocation

A second approach considers the internal allocation of resources to either photosynthetic or non-photosynthetic tissues. These tissues represent net carbon sources and sinks, respectively.

The philosophy underlying this approach is that allocation of resources to leaves (especially to leaf area) accelerates growth, whereas allocation to non-photosynthetic tissues (in this case,

roots) has an opposing effect. Allocation is an important determinant of growth rate, and this viewpoint is classically adopted by ecologists when considering the ecological strategies of plants (Grime and Hunt, 1975), resource limitations on growth (McConnaughay and Coleman, 1999; Yang and Midmore, 2005), and the growth-allocation trade-off as a constraint on life history decisions (Bazzaz *et al.*, 1987). It is also considered dynamically in relation to resource limitation in global vegetation models (e.g. Higgins and Scheiter, 2012) and in crop simulation models (e.g. Weir *et al.*, 1984; Brisson *et al.*, 1998; Jamieson *et al.*, 1998).

The change in plant mass over time is the product of leaf area ratio, net assimilation rate, and plant mass:

$$\frac{dW_P}{dt} = LAR \times NAR \times W_P \quad (5a),$$

where  $LAR$  is leaf area ratio ( $\text{m}^2$  leaf area  $\text{g}^{-1}$  plant mass) and  $NAR$  is net assimilation rate ( $\text{g carbon m}^{-2}$  leaf area  $\text{d}^{-1}$ ), remembering that carbon is equivalent to mass in these examples.

Viewed through the lens of carbon allocation, growth depends critically on the availability of photosynthetic tissue, expressed as the  $LAR$ . The  $LAR$  is in turn a product of  $SLA$ , the ratio of leaf area to leaf mass (efficiency of leaf area deployment,  $\text{m}^2 \text{g}^{-1}$  leaf mass), and  $LMR$ , the ratio of  $W_L$  to  $W_P$  (dimensionless):

$$LAR = SLA \times LMR \quad (5b),$$

Both  $SLA$  and  $LMR$  vary with  $W_P$ . At any point in time, by definition, leaf area ( $L$ , in  $\text{m}^2$ ) is therefore given by the following equation:

$$L = SLA(W_P) \times LMR(W_P) \times W_P \quad (5c),$$

where  $SLA(W_P)$  and  $LMR(W_P)$  denote  $W_P$ -dependent values of  $SLA$  and  $LMR$ . The  $LAR$  changes over time in accordance with changes in allocation during growth, and the components of  $LAR$  therefore vary with plant mass,  $W_P$ :

$$\begin{aligned} \frac{dL}{dt} = & \frac{\partial SLA}{\partial W_P} \frac{\partial W_P}{\partial t} LMR \times W_P \\ & + \frac{\partial LMR}{\partial W_P} \frac{\partial W_P}{\partial t} SLA \times W_P \\ & + \frac{\partial W_P}{\partial t} SLA \times LMR \end{aligned} \quad (5d),$$

where  $\partial SLA / \partial W_P$  and  $\partial LMR / \partial W_P$  describe the effects of allocation changing over time as plant mass changes.

The allocation perspective on growth, like the physiological perspective, can be interpreted in terms of source-sink interactions. For carbon, leaves constitute a net source while roots constitute a net sink. Thus Equation (5d) describes the change in the carbon source over time, and equivalent equations for roots would describe the change in the carbon sink.

These first two perspectives, which look at growth through the lenses of physiology and allocation, are ultimately resource-driven. The physiological perspective defines growth as being driven by carbon acquisition from, and losses to, the external environment, although in reality sink feedbacks are

also important here. The allocation perspective is driven by the allocation of carbon to structures that are responsible for its net acquisition or consumption.

### Development

The third perspective encompasses the developmental processes of organ initiation, growth, and termination. These processes represent carbon sinks.

In contrast to the first two approaches, which are resource-driven, the third perspective considers the developmental process explicitly, and this is the approach applied by developmental biologists working on growth in *Arabidopsis* and crop plants. This perspective also impinges on large-scale macroevolutionary comparisons among species, since the evolution of development must inevitably drive changes in potential growth rate, for example, in transitions between woody and herbaceous life forms (Dodd *et al.*, 1999) or in transitions between determinate and indeterminate growth (Shishkova *et al.*, 2008).

Cells divide and expand at a rate that is ultimately limited not by the speed of resource acquisition from the external environment (although this does influence meristem activity, e.g. Pritchard *et al.*, 1999; Granier *et al.*, 2007) but by intrinsic constraints set by the internal resource balance, the cell cycle and developmental programme. Again, internal source–sink interactions underpin these processes. Because cell division and expansion, and the creation of new meristems through branching constitute sinks for carbon, modelling growth from a developmental perspective places greater emphasis on the limitation of growth by sink rather than source activity. Ultimately, cell division rate is limited by molecular constraints: for example, plant genome size is negatively correlated with cell cycle time (Francis *et al.*, 2008) and with root meristem growth rate (Gruner *et al.*, 2010).

Complex formulations for organ initiation, expansion, and termination have been developed, but a simple case is considered, for illustrative purposes. If growth is considered in terms of morphogenetic constraints and development, without taking into account environmental parameters, it can be expressed as a function of the number and mass of cells:

$$\frac{dW_P}{dt} = \frac{dC}{dt} \times m \quad (6a),$$

where  $C$  is the number of cells in the plant, dependent on the division rate  $dC/dt$  in cells  $d^{-1}$ , and  $m$  is the mass of organic carbon in each cell,  $g \text{ cell}^{-1}$ .

As in Equation (4b), this can be partitioned into developmental processes occurring in leaves and in roots, where  $W_L$  and  $W_R$  refer to the dry mass of organic carbon in the leaf and root, respectively:

$$\frac{dW_P}{dt} = \frac{dW_L}{dt} + \frac{dW_R}{dt} = \frac{dC_L}{dt} \times m_L + \frac{dC_R}{dt} \times m_R \quad (6b).$$

### Unification

The three perspectives on growth can be unified to show their interrelated nature, and to illustrate the overarching

dependence of growth on source–sink relationships. While the physiological perspective focuses on metabolic processes which exchange carbon with the external environment, the allocation perspective focuses on the tissues that carry out net acquisition and drawdown of carbon, and the developmental perspective focuses on the rate of cell division in these tissues, all three perspectives are underpinned by source:sink interactions.

The mass of leaf and root tissues, seen in Equations (4b) and (6b) (relating to physiology and development, respectively), are dependent on allocation and can be expressed as follows:

$$W_L = LMR \times W_P \quad (7a),$$

$$W_R = (1 - LMR) \times W_P \quad (7b).$$

Substituting for  $dW_P/dt$  in Equation (6b) using Equation (4b) unifies the physiological and developmental perspectives:

$$\begin{aligned} & (A \times W_L) - (R_L \times W_L) - (R_R \times W_R) \\ &= \frac{dC_L}{dt} \times m_L + \frac{dC_R}{dt} \times m_R \end{aligned} \quad (8a),$$

and substituting in the definitions of  $W_L$  and  $W_R$  seen in Equations (7a) and (7b) incorporates the allocation perspective, to give:

$$\begin{aligned} & A \times LMR \times W_P - R_L \times LMR \times W_P - R_R \times (1 - LMR) \times W_P \\ &= \frac{dC_L}{dt} \times m_L + \frac{dC_R}{dt} \times m_R \end{aligned} \quad (8b),$$

where the dependence of  $SLA$  and  $LMR$  on  $W_P$  has been suppressed for ease of presentation.

This unifies the three lenses for looking at growth, and can be rearranged as:

$$\begin{aligned} A \times LMR \times W_P &= \left[ \frac{dC_L}{dt} \times m_L + R_L \times LMR \times W_P \right] \\ &+ \left[ \frac{dC_R}{dt} \times m_R + R_R \times (1 - LMR) \times W_P \right] \end{aligned} \quad (9a),$$

which is an expression of, for carbon:

$$\text{Source strength} = \text{leaf sink strength} + \text{root sink strength} \quad (9b).$$

Equation (9) illustrates an important point: it is relatively easy in a single mathematical formulation to encapsulate the intrinsic limitations on growth imposed by the physiology of resource capture, internal resource partitioning, and morphogenetic constraints on organ development. Equation (9) is not intended to be a realistic and detailed representation of growth—as discussed, it makes manifold simplifying assumptions and ignores several important components. Rather, it is intended to illustrate the potential value of taking such a unifying approach, as in the more realistic, detailed representations of the plant system developed by Chew *et al.* (2014) and Evers *et al.* (Evers *et al.*, 2010; Vos *et al.*, 2010). This unification is conceptually useful for understanding how the critical

processes of source and sink development and activity interact to limit growth in different species. A key step forwards will be model representation of the mechanisms that govern the crosstalk and interactions between different components.

Crucially, Equation (9) shows that source–sink interactions underpin all the aspects of growth described in the preceding equations. A balance between source and sink is essential for plants to grow and develop efficiently. Increased organ initiation, faster cell growth, and larger organ size will strengthen sinks; changes in the root:shoot ratio or leaf area ratio can alter the balance between carbon and nitrogen source and sink tissues; while uptake rates of carbon and mineral nutrients are primary determinants of source strength. A holistic understanding of growth rate should therefore draw on the concepts of source and sink strength, recognizing that each depends on the size and activity of the relevant tissue [Equations (1) and (2)]. Integrating molecular interactions at the tissue level (Fig. 1) with the behaviours of whole plants in terms of physiological regulation, allocation to different tissues and developmental processes will be critical for building a picture of the interactions between the three components discussed above. In order to increase crop yield effectively, it will be essential to build comprehensive growth models in which the source:sink balance is the cornerstone underpinning physiological, allocation-based, and developmental mechanisms for growth limitation. This will create an integrated perspective that allows the effects of this vital determinant of growth to be realized.

## Conclusions and recommendations

An integrated understanding of source–sink relationships, growth, and yield is a vital next step in ongoing efforts to increase crop productivity, and requires a number of key ‘unknowns’ to be addressed: (1) Which components in the plant system of sources and sinks exert the strongest control over growth in major crops? (2) How do these source and sink limitations change during the crops’ lifetimes? (3) Through what developmental or physiological mechanisms do these limitations arise? (4) Via genetic modification or selective breeding, to what extent is it possible to manipulate these processes to upregulate source and sink together, at the appropriate stage of development, to improve crop production?

We advocate the development of an integrated perspective, unifying physiological limitations on fluxes, controls on growth allocation, and the development of sink tissues, to successfully improve crop growth. A holistic view of the mechanistic interactions between sinks and sources is needed at the whole-plant scale during the trajectory of growth and development, in order to identify bottlenecks limiting growth rate. To address this knowledge gap, it will be vital to develop a greater understanding of the physiological processes operating at intermediate scales between molecular mechanisms and whole-plant traits. Ideotypes for future crops have been proposed (Sreenivasulu and Schnurbusch, 2012; Bennett *et al.*, 2012; von Caemmerer *et al.*, 2012; Reynolds *et al.*, 2012; Ort *et al.*, 2015), but reaping the maximum possible gains from

these approaches requires a parallel effort in understanding how and when source and sink capacity limit growth and yield.

## Acknowledgements

We thank Professor Andrew Fleming (Department of Animal and Plant Sciences, University of Sheffield) for helpful discussions relating to the manuscript. We are grateful to the three anonymous reviewers who provided constructive and insightful comments that significantly improved the manuscript. We thank Tiffany Bowman (Brookhaven National Laboratory) for assistance with graphic design. AW was supported by a PhD studentship from the Society for Experimental Biology (SEB). AR was supported by the United States Department of Energy contract No. DE-SC00112704 to Brookhaven National Laboratory and by the Next-Generation Ecosystem Experiments (NGEE Tropics) project that is supported by the Office of Biological and Environmental Research in the Department of Energy, Office of Science.

## References

- Acreche MM, Slafer GA.** 2009. Grain weight, radiation interception and use efficiency as affected by sink-strength in Mediterranean wheats released from 1940 to 2005. *Field Crops Research* **110**, 98–105.
- Aerts R, Chapin FS.** (2000) The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. *Adv Ecol Res* **30**, 1–67.
- Ainsworth EA, Davey PA, Hymus GJ, Osborne CP, Rogers A, Blum H, Nösberger J.** 2003. Is stimulation of leaf photosynthesis by elevated carbon dioxide concentration maintained in the long term? A test with *Lolium perenne* grown for 10 years at two nitrogen fertilization levels under Free Air CO<sub>2</sub> Enrichment (FACE). *Plant, Cell & Environment* **26**, 705–714.
- Ainsworth EA, Rogers A, Nelson R, Long SP.** 2004. Testing the ‘source–sink’ hypothesis of down-regulation of photosynthesis in elevated [CO<sub>2</sub>] in the field with single gene substitutions in *Glycine max*. *Agricultural and Forest Meteorology* **122**, 85–94.
- Ainsworth EA, Rogers A.** 2007. The response of photosynthesis and stomatal conductance to rising [CO<sub>2</sub>]: mechanisms and environmental interactions. *Plant, Cell & Environment* **30**, 258–270.
- Ainsworth EA, Leakey ADB, Ort DR, Long SP.** 2008. FACE-ing the facts: inconsistencies and interdependence among field, chamber and modeling studies of elevated [CO<sub>2</sub>] impacts on crop yield and food supply. *New Phytologist* **179**, 5–9.
- Ainsworth EA, Bush DR.** 2011. Carbohydrate export from the leaf: a highly regulated process and target to enhance photosynthesis and productivity. *Plant Physiology* **155**, 64–69.
- Álvarez F, Royo C, García del Moral LF, Villegas D.** 2008. Grain filling and dry matter translocation responses to source–sink modifications in a historical series of durum wheat. *Crop Science* **48**, 1523.
- Amthor JS.** 2000. The McCree-de Wit-Penning de Vries-Thornley respiration paradigms: 30 years later. *Annals of Botany* **86**, 1–20.
- Aranjuelo I, Sanz-Sáez A, Jauregui I, Irigoyen JJ, Araus JL, Sánchez-Díaz M, Erice G.** 2013. Harvest index, a parameter conditioning responsiveness of wheat plants to elevated CO<sub>2</sub>. *Journal of Experimental Botany* **64**, 1879–1892.
- Araújo WL, Nunes-Nesi A, Nikoloski Z, Sweetlove LJ, Fernie AR.** 2012. Metabolic control and regulation of the tricarboxylic acid cycle in photosynthetic and heterotrophic plant tissues. *Plant, Cell & Environment* **35**, 1–21.
- Arp WJ.** 1991. Effects of source-sink relations on photosynthetic acclimation to elevated CO<sub>2</sub>. *Plant, Cell & Environment* **14**, 869–875.
- Atkinson RRL, Burrell MM, Osborne CP, Rose KE, Rees M.** 2012. A non-targeted metabolomics approach to quantifying differences in root storage between fast- and slow-growing plants. *New Phytologist* **196**, 200–211.
- Badri D V, Vivanco JM.** 2009. Regulation and function of root exudates. *Plant, Cell & Environment* **32**, 666–681.
- Baraloto C, Timothy Paine CE, Poorter L, Beauchene J, Bonal D, Domenach A-M, Hérault B, Patiño S, Roggy J-C, Chave J.** 2010.

- Decoupled leaf and stem economics in rain forest trees. *Ecology Letters* **13**, 1338–1347.
- Bazzaz FA, Chiariello NR, Coley PD, Pitelka LF.** 1987. Allocating resources to reproduction and defense. *BioScience* **37**, 58–67.
- Bennett E, Roberts JA, Wagstaff C.** 2012. Manipulating resource allocation in plants. *Journal of Experimental Botany* **63**, 3391–3400.
- Bloom AJ, Chapin FS, Mooney HA.** 1985. Resource limitation in plants - an economic analogy. *Annual Review of Ecology and Systematics* **16**, 363–392.
- Brisson N, Mary B, Ripoche D, et al.** 1998. STICS : a generic model for the simulation of crops and their water and nitrogen balances. 1. Theory and parameterization applied to wheat and corn. *Agronomie* **18**, 311–346.
- Bryant J, Taylor G, Frehner M.** 1998. Photosynthetic acclimation to elevated CO<sub>2</sub> is modified by source:sink balance in three component species of chalk grassland swards grown in a free air carbon dioxide enrichment (FACE) experiment. *Plant, Cell & Environment* **21**, 159–168.
- von Caemmerer S, Farquhar GD.** 1984. Effects of partial defoliation, changes of irradiance during growth, short-term water stress and growth at enhanced p(CO<sub>2</sub>) on the photosynthetic capacity of leaves of *Phaseolus vulgaris* L. *Planta* **160**, 320–329.
- von Caemmerer S, Quick WP, Furbank RT.** 2012. The development of C<sub>4</sub> rice: current progress and future challenges. *Science* **336**, 1671–1672.
- Cameron DD, Neal AL, van Wees SCM, Ton J.** 2013. Mycorrhiza-induced resistance: more than the sum of its parts? *Trends in Plant Science* **18**, 539–545.
- Chew YH, Wenden B, Flis A, et al.** 2014. Multiscale digital *Arabidopsis* predicts individual organ and whole-organism growth. *PNAS* **111**, E4127–E4136.
- Christ RA, Körner C.** 1995. Responses of shoot and root gas exchange, leaf blade expansion and biomass production to pulses of elevated CO<sub>2</sub> in hydroponic wheat. *Journal of Experimental Botany* **46**, 1661–1667.
- Coley PD, Bryant JP, Chapin FS.** 1985. Resource availability and plant antiherbivore defence. *Science* **230**, 895–899.
- Craine JM, Lee WG, Bond WJ, Williams RJ, Johnson LC.** 2005. Environmental constraints on a global relationship among leaf and root traits of grasses. *Ecology* **86**, 12–19.
- Denison RF.** 2012. *Darwinian Agriculture: How Understanding Evolution Can Improve Agriculture*. Princeton, New Jersey: Princeton University Press.
- Díaz S, Hodgson J., Thompson K, et al.** 2004. The plant traits that drive ecosystems: Evidence from three continents. *Journal of Vegetation Science* **15**, 295–304.
- Dodd ME, Silvertown J, Chase MW.** 1999. Phylogenetic analysis of trait evolution and species diversity variation among angiosperm families. *Evolution* **53**, 732–744.
- Doehlert DC.** 1993. Sink strength: dynamic with source strength. *Plant, Cell & Environment* **16**, 1027–1028.
- Ellis RJ.** 1979. The most abundant protein in the world. *Trends in Biochemical Science* **4**, 241–244.
- Evans JR.** 1989. Photosynthesis and nitrogen relationships in leaves of C<sub>3</sub> plants. *Oecologia* **78**, 9–19.
- Eveland AL, Jackson DP.** 2012. Sugars, signalling, and plant development. *Journal of Experimental Botany* **63**, 3367–3377.
- Evers JB, Vos J, Yin X, Romero P, Van Der Putten PEL, Struik PC.** 2010. Simulation of wheat growth and development based on organ-level photosynthesis and assimilate allocation. *Journal of Experimental Botany* **61**, 2203–2216.
- Eyles A, Pinkard EA, Davies NW, Corkrey R, Churchill K, O’Grady AP, Sands P, Mohammed C.** 2013. Whole-plant- versus leaf-level regulation of photosynthetic responses after partial defoliation in *Eucalyptus globulus* saplings. *Journal of Experimental Botany* **64**, 1625–1636.
- FAO, IFAD, WFP.** 2014. *The State of Food Insecurity in the World. Strengthening the enabling environment for food security and nutrition*. Rome, FAO.
- Farage P, McKee I, Long S.** 1998. Does a low nitrogen supply necessarily lead to acclimation of photosynthesis to elevated CO<sub>2</sub>? *Plant Physiology* **118**, 573–580.
- Fatichi S, Leuzinger S, Körner C.** 2013. Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling. *New Phytologist* **201**, 1068–1095.
- Fell DA, Thomas S.** 1995. Physiological control of metabolic flux: the requirement for multisite modulation. *The Biochemical Journal* **311**, 35–39.
- Francis D, Davies MS, Barlow PW.** 2008. A strong nucleotypic effect on the cell cycle regardless of ploidy level. *Annals of Botany* **101**, 747–757.
- Freschet GT, Sward EM, Cornelissen JHC.** 2015. Integrated plant phenotypic responses to contrasting above- and below-ground resources: key roles of specific leaf area and root mass fraction. *New Phytologist* **206**, 1247–1260.
- Friedlingstein P, Meinshausen M, Arora V, Jones C, Anav A, Liddicoat S, Knutti R.** 2014. Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks. *Journal of Climate* **27**, 511–526.
- Geiger DR, Shieh W.** 1993. Sink strength: learning to measure, measuring to learn. *Plant, Cell & Environment* **16**, 1017–1018.
- Gibon Y, Pyl E, Sulpice R, Lunn JE, Höhne M, Günther M, Stitt M.** (2009) Adjustment of growth, starch turnover, protein content and central metabolism to a decrease of the carbon supply when *Arabidopsis* is grown in very short photoperiods. *Plant Cell & Environment* **32**, 859–874.
- Granier C, Cookson SJ, Tardieu F, Muller B.** 2007. Cell cycle and environmental stresses. In: Inze D, ed. *Annual Plant Reviews Volume 32: Cell Cycle Control and Plant Development*. Blackwell, 335–355.
- Grime J.** 1977. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *American Naturalist* **111**, 1169–1194.
- Grime JP, Hunt R.** 1975. Relative Growth-Rate: Its range and adaptive significance in a local flora. *Journal of Ecology* **63**, 393–422.
- Gruner A, Hoverter N, Smith T, Knight CA.** 2010. Genome size is a strong predictor of root meristem growth rate. *Journal of Botany*, Article ID 390414.
- Guilbaud CSE, Dalchau N, Purves DW, Turnbull LA.** 2015. Is ‘peak N’ key to understanding the timing of flowering in annual plants? *New Phytologist* **205**, 918–927.
- Herms D, Mattson W.** 1992. The dilemma of plants: to grow or defend. *Quarterly Review of Biology* **67**, 283–335.
- Higgins SI, Scheiter S.** 2012. Atmospheric CO<sub>2</sub> forces abrupt vegetation shifts locally, but not globally. *Nature* **488**, 209–212.
- Hubbell SP, Foster RB.** 1992. Short-term dynamics of a neotropical forest: why ecological research matters to tropical conservation and management. *Oikos* **63**, 48–61.
- IPCC.** 2014. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- Iwasa Y.** 2000. Dynamic optimization of plant growth. *Evolutionary Ecology Research* **2**, 437–455.
- Jamieson P, Semenov M, Brooking I, Francis G.** 1998. Sirius: a mechanistic model of wheat response to environmental variation. *European Journal of Agronomy* **8**, 161–179.
- Kaiser WM, Weiner H, Kandlbinder A, Tsai C-B, Rockel P, Sonoda M, Planchet E.** 2002. Modulation of nitrate reductase: some new insights, an unusual case and a potentially important side reaction. *Journal of Experimental Botany* **53**, 875–882.
- Kawakami A, Sato Y, Yoshida M.** 2008. Genetic engineering of rice capable of synthesizing fructans and enhancing chilling tolerance. *Journal of Experimental Botany* **59**, 793–802.
- Kinsman EA, Lewis C, Davies MS, Young JE, Francis D, Vilhar B, Ougham HJ.** 1997. Elevated CO<sub>2</sub> stimulates cells to divide in grass meristems: a differential effect in two natural populations of *Dactylis glomerata*. *Plant, Cell & Environment* **20**, 1309–1316.
- Klein D, Morcuende R, Stitt M, Krapp A.** 2000. Regulation of nitrate reductase expression in leaves by nitrate and nitrogen metabolism is completely overridden when sugars fall below a critical level. *Plant, Cell & Environment* **23**, 863–871.
- Klein T, Hoch G.** 2015. Tree carbon allocation dynamics determined using a carbon mass balance approach. *The New Phytologist* **205**, 147–159.

- Knorr W.** 2000. Annual and interannual CO<sub>2</sub> exchanges of the terrestrial biosphere: process-based simulations and uncertainties. *Global Ecology and Biogeography* **9**, 225–252.
- Körner C.** 1991. Some often overlooked plant characteristics as determinants of plant growth: a reconsideration. *Functional Ecology* **5**, 162–173.
- Körner C.** 2012. When growth controls photosynthesis. Paper presented to IEEE 4th International Symposium on Plant Growth Modeling, Simulation, Visualization and Applications.
- Körner C.** 2013. Growth controls photosynthesis – mostly. *Nova Acta Leopoldina* **391**, 273–283.
- Körner C, Pelaez-Riedl S, van Bel AJE.** 1995. CO<sub>2</sub> responsiveness of plants: a possible link to phloem loading. *Plant, Cell & Environment* **18**, 595–600.
- Lambers H, Freijns N, Poorter H, Hirose T, Van der Werf A.** 1989. Analyses of growth based on net assimilation rate and nitrogen productivity. Their physiological background. In: Lambers H, Cambridge M, Konings H, Pons T, eds. *Causes and Consequences of Variation in Growth Rate and Productivity of Higher Plants*. The Hague: SPB Academic Publishing.
- Lastdrager J, Hanson J, Smeekens S.** 2014. Sugar signals and the control of plant growth and development. *Journal of Experimental Botany* **65**, 799–807.
- Lawlor DW, Paul MJ.** 2014. Source/sink interactions underpin crop yield: the case for trehalose 6-phosphate/SnRK1 in improvement of wheat. *Frontiers in Plant Science* **5**, 418.
- Leakey ADB, Ainsworth EA, Bernacchi CJ, Rogers A, Long SP, Ort DR.** 2009. Elevated CO<sub>2</sub> effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* **60**, 2859–2876.
- Lekshmy S, Jain V, Khetarpal S, Pandey R, Singh R.** 2009. Effect of elevated carbon dioxide on kinetics of nitrate uptake in wheat roots. *Indian Journal of Plant Physiology* **14**, 16–22.
- Liu Y, Ahn J-E, Datta S, Salzman RA, Moon J, Huyghues-Despointes B, Pittendrigh B, Murdock LL, Koiwa H, Zhu-Salzman K.** 2005. Arabidopsis vegetative storage protein is an anti-insect acid phosphatase. *Plant Physiology* **139**, 1545–1556.
- Long SP, Ainsworth EA, Leakey ADB, Nösberger J, Ort DR.** 2006a. Food for thought: lower-than-expected crop yield stimulation with rising CO<sub>2</sub> concentrations. *Science* **312**, 1918–1921.
- Long SP, Zhu X-G, Naidu SL, Ort DR.** 2006b. Can improvement in photosynthesis increase crop yields? *Plant, Cell & Environment* **29**, 315–330.
- Lu J, Ji J.** 2006. A simulation and mechanism analysis of long-term variations at land surface over arid/semi-arid area in north China. *Journal of Geophysical Research* **111**.
- Lynch JM, Whipps JM.** 1990. Substrate flow in the rhizosphere. *Plant and Soil* **129**, 1–10.
- Marschner H.** 1995. *Mineral Nutrition of Higher Plants* (P Marschner, Ed.). London: Academic Press.
- Masle J.** 2000. The effects of elevated CO<sub>2</sub> concentrations on cell division rates, growth patterns, and blade anatomy in young wheat plants are modulated by factors related to leaf position, vernalization, and genotype. *Plant Physiology* **122**, 1399–1415.
- McConnaughay KDM, Bertson GM, Bazzaz FA.** 1993. Limitations to CO<sub>2</sub>-induced growth enhancement in pot studies. *Oecologia* **94**, 550–557.
- McConnaughay KDM, Coleman JS.** 1999. Biomass allocation in plants: ontogeny or optimality? A test along three resource gradients. *Ecology* **80**, 2581–2593.
- Metcalf CJE, Rees M, Alexander JM, Rose K.** 2006. Growth-survival trade-offs and allometries in rosette-forming perennials. *Functional Ecology* **20**, 217–225.
- Millard P.** 1988. The accumulation and storage of nitrogen by herbaceous plants. *Plant, Cell & Environment* **11**, 1–8.
- Müller-Röber B, Sonnewald U, Willmitzer L.** 1992. Inhibition of the ADP-glucose pyrophosphorylase in transgenic potatoes leads to sugar-storing tubers and influences tuber formation and expression of tuber storage protein genes. *The EMBO Journal* **11**, 1229–1238.
- Murchie EH, Pinto M, Horton P.** 2009. Agriculture and the new challenges for photosynthesis research. *New Phytologist* **181**, 532–552.
- Neales TF, Incoll LD.** 1968. The control of leaf photosynthesis rate by the level of assimilate concentration in the leaf: a review of the hypothesis. *The Botanical Review* **34**, 107–125.
- Nunes CM, O'Hara L, Primavesi L, Delatte T, Schluempmann H, Somsen G, Silva A, Fevereiro P, Wingler A, Paul MJ.** 2013. The trehalose 6-phosphate/SnRK1 signalling pathway primes growth recovery following relief of sink limitation. *Plant Physiology* **162**, 1720–1732.
- Nunes-Nesi A, Fernie AR, Stitt M.** 2010. Metabolic and signaling aspects underpinning the regulation of plant carbon nitrogen interactions. *Molecular Plant* **3**, 973–996.
- Ort DR, Merchant SS, Alric J, et al.** 2015. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences* **112**, 8529–8536.
- Pantin F, Simonneau T, Muller B.** 2012. Coming of leaf age: control of growth by hydraulics and metabolics during leaf ontogeny. *New Phytologist* **196**, 349–366.
- Pask AJD, Sylvester-Bradley R, Jamieson PD, Foulkes MJ.** 2012. Quantifying how winter wheat crops accumulate and use nitrogen reserves during growth. *Field Crops Research* **126**, 104–118.
- Patrick JW.** 1993. Sink strength: whole plant considerations. *Plant, Cell & Environment* **16**, 1019–1020.
- Paul MJ, Foyer CH.** 2001. Sink regulation of photosynthesis. *Journal of Experimental Botany* **52**, 1383–1400.
- Penning de Vries FWT.** 1975. The cost of maintenance processes in plant cells. *Annals of Botany* **39**, 77–92.
- Peterhansel C, Offermann S.** 2012. Re-engineering of carbon fixation in plants - challenges for plant biotechnology to improve yields in a high-CO<sub>2</sub> world. *Current Opinion in Biotechnology* **23**, 204–208.
- Pilkington SM, Encke B, Krohn N, Höhne M, Stitt M, Pyl E.** (2015) Relationship between starch degradation and carbon demand for maintenance and growth in *Arabidopsis thaliana* in different irradiance and temperature regimes. *Plant Cell & Environment* **38**, 157–171.
- Pollock CJ, Cairns AJ.** 1991. Fructan metabolism in grasses and cereals. *Annual Review of Plant Physiology and Plant Molecular Biology* **42**, 77–101.
- Poorter H.** 1993. Interspecific variation in the growth response of plants to an elevated ambient CO<sub>2</sub> concentration. *Vegetatio* **104**, 77–97.
- Poorter H, van der Werf A.** 1998. Is inherent variation in RGR determined by LAR at low irradiance and by NAR at high irradiance? A review of herbaceous species. In: Lambers H, Poorter H, van Vuuren M, eds. *Inherent variation in plant growth. Physiological mechanisms and ecological consequences*. Leiden: Backhuys Publishers, 309–336.
- Poorter H, Bühler J, van Dusschoten D, Climent J, Postma JA.** 2012. Pot size matters: a meta-analysis of the effects of rooting volume on plant growth. *Functional Plant Biology* **39**, 839.
- Pritchard SG, Rogers HH, Prior SA, Peterson CM.** 1999. Elevated CO<sub>2</sub> and plant structure: a review. *Global Change Biology* **5**, 807–837.
- Prusinkiewicz P, Crawford S, Smith RS, Ljung K, Bennett T, Ongaro V, Leyser O.** 2009. Control of bud activation by an auxin transport switch. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 17431–17436.
- Raines CA.** 2003. The Calvin cycle revisited. *Photosynthesis Research* **75**, 1–10.
- Ranasinghe S, Taylor G.** 1996. Mechanism for increased leaf growth in elevated CO<sub>2</sub>. *Journal of Experimental Botany* **47**, 349–358.
- Read DJ.** 1991. Mycorrhizas in ecosystems. *Experientia* **47**, 376–391.
- Reda M.** 2015. Response of nitrate reductase activity and *NIA* genes expression in roots of *Arabidopsis hxx1* mutant treated with selected carbon and nitrogen metabolites. *Plant Science* **230**, 51–58.
- Reekie EG, MacDougall G, Wong I, Hicklenton PR.** 1998. Effect of sink size on growth response to elevated atmospheric CO<sub>2</sub> within the genus *Brassica*. *Canadian Journal of Botany* **76**, 829–835.
- Reynolds M, Foulkes J, Furbank R, Griffiths S, King J, Murchie E, Parry M, Slafer G.** 2012. Achieving yield gains in wheat. *Plant, Cell & Environment* **35**, 1799–1823.
- Rogers A, Fischer B, Bryant J, Fehner M, Blum H, Raines C, Long S.** 1998. Acclimation of photosynthesis to elevated CO<sub>2</sub> under low-nitrogen nutrition is affected by the capacity for assimilate utilization.

- Perennial ryegrass under free-air CO<sub>2</sub> enrichment. *Plant Physiology* **118**, 683–689.
- Rogers A, Ainsworth EA.** 2006. The response of foliar carbohydrates to elevated [CO<sub>2</sub>]. In: Nosberger J, Long S, Blum H, Norby R, Hendrey G, Stitt M, eds. *Managed ecosystems and CO<sub>2</sub> case studies processes and perspectives*. Springer, 293–308.
- Rogers A, Ainsworth EA, Leahey ADB.** 2009. Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiology* **151**, 1009–1016.
- Rose KE, Atkinson RL, Turnbull LA, Rees M.** 2009. The costs and benefits of fast living. *Ecology Letters* **12**, 1379–1384.
- Ruuska SA, Rebetzke GJ, van Herwaarden AF, Richards RA, Fettel NA, Tabe L, Jenkins CLD.** 2006. Genotypic variation in water-soluble carbohydrate accumulation in wheat. *Functional Plant Biology* **33**, 799.
- Sage RF, Percy RW, Seemann JR.** 1987. The nitrogen use efficiency of C<sub>3</sub> and C<sub>4</sub> plants: III. Leaf nitrogen effects on the activity of carboxylating enzymes in *Chenopodium album* (L.) and *Amaranthus retroflexus* (L.). *Plant Physiology* **85**, 355–359.
- Sandve SR, Kosmala A, Rudi H, Fjellheim S, Rapacz M, Yamada T, Rognli OA.** 2011. Molecular mechanisms underlying frost tolerance in perennial grasses adapted to cold climates. *Plant Science* **180**, 69–77.
- Schnyder H.** 1993. The role of carbohydrate storage and redistribution in the source-sink relations of wheat and barley during grain filling - a review. *New Phytologist* **123**, 233–245.
- Scofield GN, Ruuska SA, Aoki N, Lewis DC, Tabe LM, Jenkins CLD.** 2009. Starch storage in the stems of wheat plants: localization and temporal changes. *Annals of Botany* **103**, 859–868.
- Sheen J.** 1990. Metabolic repression of transcription in higher plants. *The Plant Cell* **2**, 1027–1038.
- Shewry PR.** 1995. Plant storage proteins. *Biological Reviews of the Cambridge Philosophical Society* **70**, 375–426.
- Shishkova S, Rost TL, Dubrovsky JG.** 2008. Determinate root growth and meristem maintenance in angiosperms. *Annals of Botany* **101**, 319–340.
- Sitch S, Smith B, Prentice IC, et al.** 2003. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology* **9**, 161–185.
- Slewiniski TL.** 2012. Non-structural carbohydrate partitioning in grass stems: a target to increase yield stability, stress tolerance, and biofuel production. *Journal of Experimental Botany* **63**, 4647–4670.
- Smith AM, Stitt M.** 2007. Coordination of carbon supply and plant growth. *Plant, Cell & Environment* **30**, 1126–1149.
- Smith AM.** 2012. Starch in the Arabidopsis plant. *Starch/Stärke* **64**, 421–434.
- Sreenivasulu N, Schnurbusch T.** 2012. A genetic playground for enhancing grain number in cereals. *Trends in Plant Science* **17**, 91–101.
- Staswick PE.** 1990. Novel regulation of vegetative storage protein genes. *The Plant Cell* **2**, 1–6.
- Stitt M, Krapp A.** 1999. The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background. *Plant, Cell & Environment* **22**, 583–621.
- Stitt M, Zeeman SC.** 2012. Starch turnover: pathways, regulation and role in growth. *Current Opinion in Plant Biology* **15**, 282–292.
- Stokes ME, Chattopadhyay A, Wilkins O, Nambara E, Campbell MM.** 2013. Interplay between sucrose and folate modulates auxin signalling in Arabidopsis. *Plant Physiology* **162**, 1552–1565.
- Su Y-H, Liu Y-B, Zhang X-S.** 2011. Auxin-cytokinin interaction regulates meristem development. *Molecular Plant* **4**, 616–625.
- Taylor G, Ranasinghe S, Bosac C, Gardner SDL, Ferris R.** 1994. Elevated CO<sub>2</sub> and plant growth: cellular mechanisms and responses of whole plants. *Journal of Experimental Botany* **45**, 1761–1774.
- Teng S, Rognoni S, Bentsink L, Smeekens S.** 2008. The Arabidopsis *GSQ5/DOG1* Cvi allele is induced by the ABA-mediated sugar signalling pathway, and enhances sugar sensitivity by stimulating *ABI4* expression. *The Plant Journal* **55**, 372–381.
- Turnbull LA, Philipson CD, Purves DW, et al.** 2012. Plant growth rates and seed size: a re-evaluation. *Ecology* **93**, 1283–1289.
- Vos J, Evers JB, Buck-Sorlin GH, Andrieu B, Chelle M, De Visser PHB.** 2010. Functional-structural plant modelling: A new versatile tool in crop science. *Journal of Experimental Botany* **61**, 2101–2115.
- Weichert N, Saalbach I, Weichert H, et al.** 2010. Increasing sucrose uptake capacity of wheat grains stimulates storage protein synthesis. *Plant Physiology* **152**, 698–710.
- Weir AH, Bragg PL, Porter JR, Rayner JH.** 1984. A winter wheat crop simulation model without water or nutrient limitations. *Journal of Agricultural Science* **102**, 371–382.
- Woodrow IE.** 1994. Optimal acclimation of the C<sub>3</sub> photosynthetic system under enhanced CO<sub>2</sub>. *Photosynthesis Research* **39**, 401–412.
- Wright SJ, Kitajima K, Kraft NJB, et al.** 2010. Functional traits and the growth-mortality trade-off in tropical trees. *Ecology* **91**, 3664–3674.
- Yang Z, Midmore DJ.** 2005. Modelling plant resource allocation and growth partitioning in response to environmental heterogeneity. *Ecological Modelling* **181**, 59–77.
- Zaehle S, Friend AD.** 2010. Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates. *Global Biogeochemical Cycles* **24**, DOI: 10.1029/2009GB003522.
- Zeeman SC, Kossmann J, Smith AM.** 2010. Starch: its metabolism, evolution, and biotechnological modification in plants. *Annual Review of Plant Biology* **61**, 209–234.
- Zhu X-G, de Sturler E, Long SP.** 2007. Optimizing the distribution of resources between enzymes of carbon metabolism can dramatically increase photosynthetic rate: a numerical simulation using an evolutionary algorithm. *Plant Physiology* **145**, 513–526.
- Zuther E, Hoermiller II, Heyer AG.** 2011. Evidence against sink limitation by the sucrose-to-starch route in potato plants expressing fructosyltransferases. *Physiologia Plantarum* **143**, 115–125.