

DOES FOSSIL FUEL COMBUSTION LEAD TO GLOBAL WARMING?

Stephen E. Schwartz
Environmental Chemistry Division
Brookhaven National Laboratory
Upton, NY 11973

January 1993

By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

This research was performed under the auspices of the United States Department of Energy, under Contract No. DE-AC02-76CH00016.

DOES FOSSIL FUEL COMBUSTION LEAD TO GLOBAL WARMING?

Stephen E. Schwartz
Environmental Chemistry Division
Brookhaven National Laboratory
Upton NY 11973 USA

[Abstract]

Tropospheric sulfate aerosols produced by atmospheric oxidation of SO₂ emitted from fossil fuel combustion scatter solar radiation and enhance the reflectivity of clouds. Both effects decrease the absorption of solar radiation by the earth-atmosphere system. This cooling influence tends to offset the warming influence resulting from increased absorption of terrestrial infrared radiation by increased atmospheric concentrations of CO₂. The sulfate forcing is estimated to be offsetting 70% of the forcing by CO₂ derived from fossil fuel combustion, although the uncertainty of this estimate is quite large--range 28 to 140%, the latter figure indicating that the present combined forcing is net cooling. Because of the vastly different atmospheric residence times of sulfate aerosol (about a week) and CO₂ (about 100 years), the cooling influence of sulfate aerosol is exerted immediately, whereas most of the warming influence of CO₂ is exerted over more than 100 years. Consequently the total forcing integrated over the entire time the materials reside in the atmosphere is net warming, with the total CO₂ forcing estimated to exceed the sulfate forcing by a factor of 4 (uncertainty range 2 to more than 10). The present situation in which the forcing by sulfate is comparable to that by CO₂ is shown to be a consequence of the steeply increasing rates of emissions over the industrial era.

INTRODUCTION

It is generally accepted that fossil fuel combustion is responsible for much of the increase in atmospheric carbon dioxide concentrations above preindustrial values and that this increased CO₂ results in an increased longwave (infrared) radiative forcing of the troposphere that leads to a warming of the earth's surface.{Houghton, 1990 #1} Recently attention has been called to a shortwave radiative forcing by tropospheric sulfate aerosols produced by atmospheric oxidation of SO₂ emitted from fossil fuel combustion.{Twomey, 1984 #2;

Schwartz, 1988 #3; Wigley, 1992 #4; Wigley, 1991 #5; Kaufman, 1991 #6; Charlson, 1991 #7; Twomey, 1991 #8; Charlson, 1992 #9} These aerosols scatter solar radiation, increasing clear-sky planetary reflectivity. Increased concentrations of aerosol particles also lead to increased reflectivity of clouds. Both effects are thought to contribute a cooling influence on the planetary energy balance that to some extent offsets the warming influence due to increased CO₂. Accurate knowledge of the net forcing due to fossil fuel combustion is necessary both for interpretation of temperature trends over the industrial era and for formulation of future policies regarding fossil fuel combustion.{Houghton, 1990 #1; Houghton, 1992 #10; Wigley, 1992 #4; Schlesinger, 1992 #11; Kelly, 1992 #12} It is therefore necessary to examine the climate influence of fossil fuel combustion taking into account both the infrared forcing due to increased concentrations of CO₂ and the shortwave forcing due to sulfate aerosol. A major difference in the two forcings is the residence time of the materials in the atmosphere: decades to centuries for CO₂, days to weeks for sulfate. Another key parameter governing the relative magnitudes of these forcings is the sulfur content in the fuel, which has decreased in recent years and which may be expected to decrease further in response to controls on SO₂ emissions aimed at reducing acid deposition.

In this article I first describe and compare the two modes of radiative forcing and review estimates of the normalized instantaneous radiative forcing due to each of the two substances. I then examine the extent to which the relative magnitudes of warming and cooling depend on the time history of emissions, in view of the great differences in atmospheric residence times of sulfate and CO₂. I show that as a consequence of the greatly differing residence times, the net radiative forcing by the two substances, emitted in constant ratio, can be quite different for different time profiles of emissions. I then apply this analysis to a time history of fossil fuel combustion over the industrial period to assess the relative influence of these two opposing forcings up to the present time and consider implications for the future.

Comparison of present radiative influences of fossil-fuel derived CO₂ and sulfate

It is well established that the mixing ratio of atmospheric CO₂ has been increasing substantially over the industrial era{Watson, 1990 #13} from the preindustrial value of 280 ppm (parts per million by volume) to the present (1990) value of 353 ppm. The radiative influence of this excess CO₂ is quantified as radiative forcing of the earth-troposphere system, which is estimated, again for 1990, as 1.5 W m⁻²;
{Shine, 1990 #14} this is the greatest single contribution to anthropogenic greenhouse gas forcing, which is estimated as 2.45 W m⁻² in the

aggregate. Although the increase in CO₂ concentration is well established, apportionment of that increase to fossil fuel combustion and to other activities such as deforestation and land use changes is less certain. The contribution of fossil fuel combustion to the net increase in CO₂ flux into the atmosphere over the industrial period as a whole is estimated as $63 \pm 10\%$;{Shine, 1990 #14} this relative contribution has been increasing over this period, with the present (1980-1989) contribution estimated as $76 \pm 12\%$. In view of the long residence time of excess CO₂ in the atmosphere (decades to centuries{Shine, 1990 #14}), it is not clear a priori what are the appropriate values to employ in estimating the fossil-fuel contribution to excess CO₂ forcing, so I take this range as $71 \pm 17 \%$; the radiative forcing attributed to fossil fuel CO₂ is thus estimated as $1.07 \pm 0.26 \text{ W m}^{-2}$.

The radiative forcing due to anthropogenic sulfate has recently been the subject of much scrutiny. The steady-state global-mean direct radiative forcing was estimated by a box model calculation{Charlson, 1992 #9} as -1.3 W m^{-2} , uncertain to a factor of 2, the principal contributions to the uncertainty being the fractional conversion of SO₂ to sulfate and the mean atmospheric residence time of the sulfate aerosol, which is about a week. (The negative sign indicates that this radiative influence is one of cooling, in contrast to the warming influence of excess CO₂). Calculations using sulfate concentration fields obtained with a three dimensional global model{Langner, 1991 #15} led to an estimate of global mean forcing of -0.60 W m^{-2} ,{Charlson, 1991 #7} again uncertain to a factor of 2. In addition to the direct forcing Charlson et al.{Charlson, 1992 #9} estimated the so-called indirect forcing due to enhancement of cloud albedo, based on an assumed 15% global-mean increase in the concentration of droplets in marine stratus clouds, that in turn is based on a comparison of measured sulfate aerosol concentrations at locations remote from industrial emissions in the Northern and Southern Hemispheres.{Schwartz, 1988 #3} The resulting global-mean cloud forcing was estimated as -1 W m^{-2} , additive to the direct forcing; however, this estimate must be viewed as even more uncertain than that of the direct forcing because of the highly nonuniform spatial distribution of the aerosol, the unknown dependence of cloud droplet concentration on sulfate concentration, and the nonlinear dependence of cloud reflectance on cloud droplet concentration. Recently Wigley and Raper{Wigley, 1992 #4} reviewing the available evidence on sulfate forcing estimated the total forcing (direct plus cloud) as $-0.75 \pm 0.38 \text{ W m}^{-2}$. Since the focus of the present article is restricted to the effects of fossil fuel combustion, the above estimates should be reduced by 8% to account for non fossil fuel anthropogenic sulfur

emissions, largely from smelting of non-ferrous metals;{Spiro, 1992} with this reduction the Wigley-Raper{Wigley, 1992 #4} estimate for the radiative forcing by fossil-fuel sulfate becomes $-0.70 \pm 0.35 \text{ W m}^{-2}$.

The several estimates of sulfate forcing range widely. They can be refined only by improved descriptions of the distribution and radiative properties of this aerosol. The sulfate aerosol forcing is distributed quite differently in space and time from the greenhouse gas forcing and consequently equivalent climate response is not necessarily expected. Nonetheless it is clear that this forcing is substantial as measured by the yardstick of fossil fuel CO_2 forcing or total greenhouse gas forcing. If we take the Wigley-Raper{Wigley, 1992 #4} estimate as a conservative recent estimate of the sulfate forcing, and combine the range of this estimate with the range of estimates of fossil fuel CO_2 forcing given above, then we obtain an estimate for the present net radiative forcing due to fossil fuel combustion of $0.37 \pm 0.61 \text{ W m}^{-2}$, i.e. ranging from $+0.98 \text{ W m}^{-2}$ (net warming) to -0.24 W m^{-2} (net cooling).

In view of the substantial uncertainty, even in the sign of the net forcing, it appears that a systematic comparison of the forcings of CO_2 and sulfate is warranted. Further, in view of the widely differing residence times of the two substances in the atmosphere, such a comparison must focus on the time dependence of the forcings by these materials subsequent to their introduction into the atmosphere.

Global warming potentials and integrals

To compare the radiative forcing of sulfate aerosols to that of CO_2 , I employ and extend the global warming potential (GWP) approach,{Shine, 1990 #14; Fisher, 1990 #17} which was developed to facilitate comparison of the climate influence of different greenhouse gases. The GWP is a single index that accounts for the climate influence that would result from introduction of a specified incremental amount of an infrared-active gas into the atmosphere, relative to that for carbon dioxide. As generally employed{Shine, 1990 #14} the GWP of a substance i is a dimensionless quantity evaluated as the time integrated radiative forcing of the earth-troposphere system due to introduction of a unit incremental amount of the substance into the atmosphere, ratioed to the same quantity for CO_2 :

$$\text{GWP}_i(T) \equiv \int_0^T a_i f_i(t) dt / \int_0^T a_{\text{CO}_2} f_{\text{CO}_2}(t) dt \quad (1)$$

Here a_i represents the instantaneous radiative forcing due to an incremental amount (1 kg) of substance i in the atmosphere, and $f_i(t)$, the decay profile of the substance, is the fraction of material remaining in the atmosphere at a time t subsequent to its introduction. The integral is taken over a period of time T that is denoted the "time horizon" of concern; if the substance has a different decay profile in the atmosphere from that of CO_2 , the GWP will depend on the time horizon that is employed. This useful quantity encompasses the effects of both the magnitude of instantaneous radiative forcing and residence time of the substance in describing its climatic influence. Although criticized as not accurately representing differences in the details of forcings by different greenhouse gases, {Wang, 1991 #18} the approach continues to be employed {Houghton, 1992 #10} as a useful first-order estimator of total global influences.

It is seen that the GWP of a particular substance i depends not only on the radiative and persistence properties of that substance, but also on the corresponding properties of CO_2 . In view of the vastly differing time scales characterizing the removal of CO_2 and aerosol sulfate from the atmosphere, the two integrals comprising the numerator and denominator of the GWP exhibit quite different time behavior. Consequently much more insight can be gained by examining the two integrals individually than can be gained from examination only of the ratio. Moreover uncertainties in the properties of both the substance and CO_2 contribute to uncertainties in the GWP. As it turns out there are substantial uncertainties in the atmospheric decay profile not only of sulfate aerosol but also of excess CO_2 . Indeed, the Intergovernmental Panel on Climate Change (IPCC) states that because of uncertainties associated with specifying the lifetime of CO_2 , "presentation of the GWP relative to CO_2 may not be the ideal choice." {Shine, 1990 #14} For these reasons I introduce a quantity that I denote as the Greenhouse Warming Integral (GWI):

$$\text{GWI}_i(T) \equiv \int_0^T a_i f_i(t) dt \quad (2)$$

Unlike the GWP, the dependence of the GWI on the time horizon T depends on the residence time profile $f_i(t)$ of only a single substance. In contrast to the GWP, the GWI has dimension; for a_i expressed in units $\text{W m}^{-2} \text{kg}^{-1}$, and for time taken in seconds, the GWP has units $\text{J m}^{-2} \text{kg}^{-1}$. In the limit $T \rightarrow \infty$ the GWI admits to a ready interpretation: it is the global-mean incremental heating per unit surface area of the earth resulting from the introduction of 1 kg of the substance i into the atmosphere. If the forcing per kilogram a_i is a constant this total

GWI is simply expressed as the product of a_i times the mean residence time of the material in the atmosphere τ_i :

$$\text{GWI}_i^\infty = a_i \int_0^\infty f_i(t) dt = a_i \tau_i \quad (3)$$

where τ_i is the mean residence time of the material in the atmosphere. For a gas whose decay profile is exponential, the partial GWI for time horizon T is $a_i \tau_i [1 - f_i(T)] = a_i \tau_i [1 - \exp(-T / \tau_i)]$.

For a species whose residence time is sufficiently short relative to secular changes in source strength that it may be considered to be in steady state, the total GWI may equivalently be calculated as the global-mean incremental radiative forcing F_i (W m^{-2}) divided by the total global source strength Q_i (kg s^{-1}) of the substance. [Schwartz, 1979 #19]

$$\text{GWI}_i^\infty = F_i / Q_i \quad (4)$$

The incremental forcing per kilogram of material emitted into the atmosphere can thus be evaluated from (3) as

$$a_i = F_i / Q_i \tau_i \quad (5)$$

The GWP approach has previously been applied only to greenhouse gases, and in its 1992 update the IPCC explicitly indicated that no attempt was made to define a GWP for anthropogenic sulfate emissions, in view of "the very different character of the forcing due to aerosol sulphate as compared to that of the greenhouse gases." [Isaksen, 1992 #20] In fact however the approach is readily extended to aerosol sulfate by means of a sign convention such that a negative GWP or GWI denotes a cooling influence. The extension appears to be useful, and I make use of it here in order to compare the magnitude of radiative forcing of climate by fossil-fuel CO_2 and sulfate aerosol.

Global warming integral for sulfate aerosol

To derive a GWI for sulfate I use the radiative forcing estimates noted above together with the corresponding emissions of anthropogenic SO_2 . Scattering of shortwave radiation (direct forcing) is to good approximation linear in emitted SO_2 (the extent of SO_2 oxidation is assumed linear in SO_2 concentration, the resulting sulfate aerosol is optically thin, and the removal of the aerosol from the atmosphere, mainly by precipitation, is assumed independent

of the aerosol concentration). The Charlson et al. {Charlson, 1992 #9} box model calculation yielded, for annual anthropogenic emissions of SO_2 , Q_S , taken as 90 Tg S yr^{-1} and sulfate mean residence time of 7 days, a steady-state global-mean direct radiative forcing of -1.3 W m^{-2} . The three dimensional global model of Langner and Rodhe {Langner, 1991 #15} led to an estimate {Charlson, 1991 #7} of the steady-state global-mean direct radiative forcing of -0.60 W m^{-2} , for the SO_2 emission rate taken as 71 Tg S yr^{-1} ; in these calculations the mean residence time of sulfate aerosol was found to be about 5 days. From these estimates and equations (4) and (5) it is possible to evaluate the total (infinite time) GWI of tropospheric sulfate aerosol, and the instantaneous average forcing per kilogram emitted sulfur, a_S , as given in Table 1.

Several points should be noted regarding this table. First, the GWI is negative, indicative of a cooling influence. Second, the factor of two difference in forcing estimated by the two models is seen to be due in large part to the different emission rates and mean residence times employed; the actual forcing per kilogram of sulfur emitted into the atmosphere, a_S , differs between the models by only about 20%. However this agreement should not necessarily be taken as indicative of the certainty of the estimates. The estimates of GWI_S and a_S should be viewed as uncertain to roughly a factor of two. For the purpose of the present comparison of radiative forcing of sulfate and CO_2 from fossil fuel combustion I employ the values $360 \mu\text{J m}^{-2} \text{ kg}^{-1}$ and $690 \text{ pW m}^{-2} \text{ kg}^{-1}$, respectively, corresponding to a mean residence time of sulfate of 6 days. These values, together with the present (1989) rate of emissions of SO_2 from fossil fuel combustion, (87 Tg S yr^{-1} , as inferred from the present rate of CO_2 emissions, 5.8 Pg C yr^{-1} , and the present sulfur to carbon ratio in fossil fuel, 0.014, Figure 1) yield for the present global average direct forcing by sulfate aerosol -1.0 W m^{-2} , essentially the same magnitude as the estimate given above for present radiative forcing by fossil-fuel derived CO_2 . As improved estimates of the several quantities become available, the GWI estimates obtained here may be scaled appropriately.

In addition to the direct forcing there is also thought to be a contribution to the GWI from the cloud forcing. Charlson et al. {Charlson, 1992 #9} estimate this forcing as -1 W m^{-2} , but highly uncertain. For reference the corresponding contributions to GWI and a_S are also given in Table 1. However in view of the uncertainty in estimating the cloud forcing, I use only the direct forcing due to clear-sky light scattering in comparisons with the CO_2 forcing; the resulting comparisons can be scaled appropriately to include the cloud forcing.

Global warming integral for CO₂

The incremental longwave radiative forcing of the troposphere by CO₂ is approximately logarithmic in CO₂ concentration, such that a doubling of CO₂ concentration corresponds to an incremental forcing of approximately 4 W m⁻². Thus the instantaneous forcing of an incremental amount of CO₂ added to the atmosphere $\Delta[\text{CO}_2]$ can be approximated as:{Shine, 1990 #14}

$$a_{\text{CO}_2} = \left(\frac{4 \text{ W m}^{-2}}{\ln 2} \right) \Delta \ln[\text{CO}_2] = \left(5.8 \text{ W m}^{-2} \right) \frac{\Delta[\text{CO}_2]}{[\text{CO}_2]} \quad (6)$$

where $[\text{CO}_2]$ is the amount initially present in the atmosphere. For $[\text{CO}_2]$ taken as 670 Pg C, equivalent to 315 ppm, intermediate between preindustrial and present values, $a_{\text{CO}_2} = 8.6 \times 10^{-15} \text{ W m}^{-2} \text{ per kg C}$. Because the departure from linearity over this range of $[\text{CO}_2]$ is fairly small, the GWI of CO₂ is adequately approximated as $a_{\text{CO}_2} \int_0^T f_{\text{CO}_2}(t) dt$.

In contrast to sulfate, excess CO₂ from fossil fuel combustion remains in the atmosphere for decades to centuries. The decay profile of excess CO₂ in the atmosphere is not well established. Multi-compartment model calculations{Maier-Reimer, 1987 #21; Siegenthaler, 1983 #22} indicate a nonexponential decay, with an initial half-life of about 50 yr but with the second half-life much greater, about 250 yr, Figure 2; in fact according to such models the incremental CO₂ mixing ratio may never return to zero, depending on assumptions of the ultimate geochemical fate of incremental CO₂ added to the atmosphere. Such a profile, for which the equivalent mean residence time was stated to be approximately 120 years, was employed by the IPCC{Houghton, 1990 #1; Houghton, 1992 #10} in GWP estimates.

Alternative estimates of the mean residence time of CO₂ in the atmosphere can be inferred from the time dependence of CO₂ concentration in the atmosphere in excess of the preindustrial value, the rate of increase in atmospheric CO₂, and the rate of CO₂ emissions from industrial activity as a function of time. Using the data given by IPCC for these quantities I infer the mean residence time of excess CO₂ in the present atmosphere to be about 85 years. A value of 68 years can be inferred from estimates of carbon fluxes to surface ocean water derived from the rate of decrease of ¹³C concentration.{Quay, 1992 #23} In view of this range of values I take the mean residence time as 100 years, uncertain to 30%; the corresponding exponential decay profiles are given in Figure 2. It is seen that initially these

profiles initially decay more slowly than the nonlinear model, but that ultimately they decay much more rapidly. The total GWI (time horizon $T \rightarrow \infty$) is quite sensitive to the long-time behavior of these profiles.

We are now prepared to evaluate GWI(T) for CO₂. Figure 3 shows this quantity for the three exponential decay profiles ($\tau_{\text{CO}_2} = 70, 100, 130$ years) and for the nonexponential profile of Maier-Reimer and Hasselmann.{Maier-Reimer, 1987 #21} As expected these profiles increase from zero over a time scale of some 200 years. The final value for the exponential profiles is $\text{GWI}_C^\infty = (27 \pm 8) \mu \text{J m}^{-2}$ per kg C emitted into the atmosphere as CO₂; for the nonexponential decay profile the value is greater and in fact may continue to increase without bound, depending on assumptions regarding the ultimate fate of incremental CO₂.

Because time scale of the decay of excess CO₂ in the atmosphere is long compared to that characterizing the increase in the rate of emission of CO₂ over the industrial period, (Fig. 1), atmospheric CO₂ cannot be considered to be in steady state; indeed the annual increase in the burden of atmospheric CO₂ is a substantial fraction of the annual fossil fuel emissions. For this reason it is mandatory to consider the time-dependence of emission and forcing in comparing the climate forcing of sulfate aerosol and CO₂. In the long run the relative contributions of the two substances can be compared by consideration of the infinite-time integrals. However on a time scale over which CO₂ is not in steady state it is necessary to employ the partial GWI for CO₂ and to take explicit account of the historical time profile of CO₂ emissions. As shown below, this is crucial for comparison of the two forcings over the industrial period.

Comparison of infinite time GWI's of CO₂ and sulfate aerosol

With the total GWI's for CO₂ and sulfate it is possible to compare the net radiative forcing of the two species, that is the net GWI for fossil fuel combustion, integrated over the entire residence times of the two materials in the atmosphere. Here I choose to express this forcing in terms of the unit mass emissions of carbon, expressing the emission of sulfur in terms of the mass ratio in emissions (Q_S/Q_C). Thus the net GWI for fossil fuel combustion normalized to emission of carbon is

$$\text{GWI}_{\text{ff}}^\infty = \text{GWI}_C^\infty + (Q_S / Q_C) \text{GWI}_S^\infty \quad (7a)$$

For the two GWI's evaluated by (3) we have

$$\text{GWI}_{\text{ff}}^{\infty} = a_{\text{CO}_2} \tau_{\text{CO}_2} + (Q_S / Q_C) a_S \tau_{\text{SO}_4^{2-}} \quad (7b)$$

Equation 7 permits determination of a critical ratio of emissions of SO₂ and CO₂, $(Q_S / Q_C)^*$, such that the incremental forcings negate each other, i.e., such that $\text{GWI}_{\text{ff}}^{\infty} = 0$, as

$$(Q_S / Q_C)^* = -\frac{\text{GWI}_C^{\infty}}{\text{GWI}_S^{\infty}} \quad (8)$$

Or by (7b)

$$(Q_S / Q_C)^* = -\frac{a_{\text{CO}_2} \tau_{\text{CO}_2}}{a_S \tau_{\text{SO}_4^{2-}}} \quad (8b)$$

For $(Q_S / Q_C) < (Q_S / Q_C)^*$ the net forcing is positive, i.e., warming, whereas for $(Q_S / Q_C) > (Q_S / Q_C)^*$ the net forcing is negative. For the total GWI's obtained above $(Q_S / Q_C)^*$ is equal to 0.075, with an uncertainty, due largely to the uncertainty in the GWI of sulfate, of about a factor of 2. This value is roughly four times the mass ratio of sulfur to carbon in emissions from fossil fuel combustion (Figure 1); the emission-weighted average over the period 1860 to 1987, $\overline{(Q_S / Q_C)}$, is 0.019. Consequently when the CO₂ forcing over its entire residence time in the atmosphere is considered, the positive forcing due to CO₂ exceeds the negative forcing due to sulfate aerosol, and that the net effect of fossil fuel combustion, evaluated for the total period of time over which the excess CO₂ remains in the atmosphere, is unequivocally one of heating. This result would also apply to the steady-state emissions scenario, that is, to a situation in which both substances are emitted into the atmosphere at constant rate for a period long compared to the mean residence time of excess CO₂. It is clear, however, that the conclusion might not be the same with respect to the actual net forcing experienced to date over the industrial era, especially as the rate of emissions has been increasing strongly over this period, and therefore much of the ultimate heating influence of excess CO₂ still present in the atmosphere has not yet been exerted.

Model for comparing CO₂ and sulfate forcing with exponentially increasing emissions

It should be apparent by now that the net radiative forcing experienced by the earth-troposphere system due to CO₂ and sulfate aerosol resulting from fossil fuel combustion must be considered as a function of time following emission. Here I explore the implications of a

CO₂ emission profile that is increasing rapidly relative to the time scale of the decay profile of excess CO₂ in the atmosphere. By considering an exponential growth profile for emissions and an exponential decay profile for atmospheric CO₂ one obtains a great simplification in the algebra that leads to insight. We have already seen that the decay profile of CO₂, f_{CO_2} , appears to be fairly accurately represented by an exponential, with e-folding time of about 100 years. Figure 4 shows that the time history of emissions of fossil fuel CO₂ over the industrial period is fairly accurately represented by exponential growth with an e-folding time of 30 years.

I now evaluate the radiative forcing of a species i that is emitted at an exponentially increasing rate $Q_i(t') = Q_i(t_0)\exp[(t' - t_0)/\tau_e]$ and exhibits an exponential decay profile $f_i(t) = \exp[-(t - t')/\tau_i]$. The radiative forcing at time t is given by

$$\begin{aligned} F_i(t) &= a_i \int_{-\infty}^t Q_i(t') f_i(t - t') dt' \\ &= a_i Q_i(t) \left(\frac{\tau_i \tau_e}{\tau_i + \tau_e} \right) \end{aligned} \quad (9)$$

It is seen that the forcing exerted at any given time t is in constant proportion to the emission rate at that time, and thus by analogy to equation (4) I define an effective GWI for this situation as

$$\text{GWI}_i^{\text{exp}} \equiv F_i(t) / Q_i(t) = a_i \left(\frac{\tau_i \tau_e}{\tau_i + \tau_e} \right) \quad (10)$$

This expression differs from that for GWI_i^{∞} in that the mean atmospheric residence time τ_i is replaced by the quantity in brackets, the harmonic sum of τ_i and the e-folding time of emissions τ_e . For sulfate aerosol, for which $\tau_{\text{SO}_4^{2-}} \ll \tau_e$ this quantity equals $\tau_{\text{SO}_4^{2-}}$ and

$$\text{GWI}_S^{\text{exp}} = a_S \tau_{\text{SO}_4^{2-}}, \quad (11a)$$

that is, $\text{GWI}_S^{\text{exp}}$ is identical to GWI_S^{∞} . For CO₂ the situation is nearly reversed, i.e. the atmospheric residence time of excess CO₂, τ_{CO_2} , is almost sufficiently greater than τ_e that the latter can be neglected in the summation in the denominator of (10). If that were the case we would have

$$\text{GWI}_C^{\text{exp}} = a_{\text{CO}_2} \tau_e \quad (11b)$$

In fact that is not strictly accurate so equation (11b) must be replaced by

$$\text{GWI}_C^{\text{exp}} = a_{\text{CO}_2} \tau_e (1 + \tau_e / \tau_i)^{-1} \quad (11c)$$

where for $\tau_e = 30$ years and $\tau_{\text{CO}_2} = 100$ years the correction factor is about 0.77, yielding a value of about 23 years. It is very insightful to compare equation (11) with equation (3). In a situation of exponentially increasing emissions, with e-folding time associated with the increase of emissions τ_e , for the long lived CO_2 the mean atmospheric residence time τ_{CO_2} has effectively been replaced by the considerably smaller τ_e (in fact, by the even slightly smaller quantity, $\tau_e (1 + \tau_e / \tau_{\text{CO}_2})^{-1}$). For $\tau_{\text{CO}_2} = 100 \pm 30$ years, this has the effect of reducing GWI_C from 27 ± 8 to $6.2 \pm 0.5 \mu\text{J m}^{-2}$ per kg C. As may be seen from Figure 3, this value corresponds to the $\text{GWI}(T)$ for a value of T of about 23 years.

In contrast to the situation for CO_2 , for the short-lived sulfate there is no difference between the GWC evaluated for the exponentially increasing emissions and the total GWC. This implies that for a given emission ratio (Q_S / Q_C) the magnitude of the warming effect of CO_2 relative to the cooling effect of sulfate is substantially reduced from that which would be experienced over the entire residence times of the materials, as evaluated above. The net time-dependent forcing due to fossil fuel combustion can be evaluated as

$$\begin{aligned} F_{\text{ff}}(t) &= F_C(t) + F_S(t) \\ &= Q_C(t) \left[\text{GWI}_C^{\text{exp}} + (Q_S / Q_C) \text{GWI}_S^{\text{exp}} \right] \\ &= Q_C(t) \left[a_{\text{CO}_2} \tau_e (1 + \tau_e / \tau_{\text{CO}_2})^{-1} + (Q_S / Q_C) a_S \tau_{\text{SO}_4^{2-}} \right] \\ &\approx Q_C(t) \left[a_{\text{CO}_2} \tau_e + (Q_S / Q_C) a_S \tau_{\text{SO}_4^{2-}} \right] \end{aligned} \quad (12)$$

where as above Q_S is expressed in terms of the mass ratio in emissions (Q_S / Q_C). Also as above we evaluate a critical emissions ratio such that the net forcing is zero (cf. equation 8):

$$(Q_S / Q_C)^* = - \frac{\text{GWI}_C^{\text{exp}}}{\text{GWI}_S^{\text{exp}}} = - \frac{a_{\text{CO}_2} \tau_e (1 + \tau_e / \tau_{\text{CO}_2})^{-1}}{a_S \tau_{\text{SO}_4^{2-}}} \approx - \frac{a_{\text{CO}_2} \tau_e}{a_S \tau_{\text{SO}_4^{2-}}} \quad (13)$$

As a consequence of the change in the effective GWI for CO₂, the value for $(Q_S/Q_C)^*$ of 0.075 obtained above with the total GWI's is decreased, for $\tau_{\text{CO}_2} = 100$ years, by a factor of 0.23, i.e., to 0.017, again with an uncertainty estimate of a factor of 2. This value is essentially the same as the emission-weighted average $(\overline{Q_S}/\overline{Q_C})$ over the period 1860 to 1987, 0.019. To the extent that this model is applicable, this calculation suggests that the net radiative forcing by fossil fuel combustion emissions, at any time during the exponential growth period, is near zero.

This analysis shows that for an exponentially increasing emissions profile, similar to that actually experienced over the industrial era, the net global warming influence of the two species, having two quite different atmospheric residence times, is quite different from that which would be expected on the basis of the infinite-time global warming integrals. The other side of the coin, however, is the implication that for the values $\tau_{\text{CO}_2} = 100$ years and $\tau_e = 30$ years employed in the calculation, only 23% of the total GWI of all the excess CO₂ introduced into the atmosphere according to this emissions profile prior to a given time has been exerted up until that time; the remainder of the GWI of the CO₂ is yet to be exerted, whereas the total GWI for the sulfate is exerted essentially immediately. The fact that the warming influence of CO₂ is to great extent offset by the cooling influence of sulfate, despite the much greater warming influence of CO₂ than cooling influence of sulfate inferred from the total GWI's, is a subtle consequence of the exponentially increasing emissions profile, i.e., the warming influence of past CO₂ emissions is offset largely by current SO₂ emissions. Such a situation can be maintained indefinitely, but only provided that the emissions of both substances continue to increase at the same exponential rate. However such a pattern of emissions implies that the burden of unexpressed CO₂ forcing also is continuing to increase exponentially. And of course exponential growth rates are hard to sustain indefinitely.

Comparison of sulfate and CO₂ forcing based on emissions over industrial era

I now extend the analysis given above to consideration of the forcing resulting from the actual profile of CO₂ and SO₂ emissions from fossil fuel combustion over the industrial era. This analysis permits comparison of the carbon and sulfur forcings that have occurred only over this period, i.e., excluding the subsequent influence of CO₂ emitted during this period and still remaining in the atmosphere.

From equation (3) it may be seen that the integrated forcing exerted over the time period extending from the time of emission t_e to a subsequent time T is given by

$$\text{IF}(T, t_e) = Q_i(t_e) \int_{t_e}^T a_i f_i(t) dt = Q_i(t_e) \text{GWI}_i(T, t_e) \quad (14)$$

Consequently the total forcing exerted over a time period t_0 to T , representing the time period from the beginning of industrialization to the present, is given by

$$\text{IF}(T) \equiv \int_{t_0}^T \text{IF}(T, t_e) dt_e = \int_{t_0}^T Q_i(t_e) \text{GWI}_i(T, t_e) dt_e, \quad (15)$$

that is, by the integral over time of the GWI weighted by the emission rate. In the case of sulfate the GWI reaches its final and maximum value GWI_S^∞ essentially instantaneously, so that the integrated forcing is equal to the product of GWI_S^∞ times the total SO_2 emissions over the period.

$$\text{IF}_S(T) = \text{GWI}_S^\infty \int_{t_0}^T Q_S(t_e) dt_e \quad (16)$$

In turn this can be expressed in terms of the mean emission ratio $\overline{(Q_S / Q_C)}$ as

$$\text{IF}_S(T) = \text{GWI}_S^\infty \overline{(Q_S / Q_C)} E_C^{\text{tot}} \quad (17)$$

where E_C^{tot} is the total emission of CO_2 over the time period,

$$E_C^{\text{tot}} = \int_{t_0}^T Q_C(t_e) dt_e \quad (18)$$

In the case of CO_2 the integral in (15) must be evaluated for the historical emission rate and an assumed decay profile. It can be expressed as the product of the total emission of CO_2 times the emission weighted mean GWI, $\langle \text{GWI}_C \rangle$:

$$\text{IF}_C(T) = E_C^{\text{tot}} \langle \text{GWI}_C \rangle \quad (19)$$

where

$$\langle \text{GWI}_C \rangle = \int_{t_0}^T \text{GWI}_C(T, t_e) Q_C(t_e) dt_e / \int_{t_0}^T Q_C(t_e) dt_e \quad (20)$$

The net integrated forcing due to fossil fuel combustion over the time period t_0 to T and exerted over this period is

$$\begin{aligned} \text{IF}_{\text{ff}}(T) &= \text{IF}_C(T) + \text{IF}_S(T) \\ &= E_C^{\text{tot}} \left(\langle \text{GWI}_C \rangle + \overline{(Q_S / Q_C)} \text{GWI}_S^\infty \right) \end{aligned} \quad (21)$$

As above we define a critical emissions ratio such that the net forcing is zero:

$$(Q_S / Q_C)^* = - \frac{\langle \text{GWI}_C \rangle}{\text{GWI}_S^\infty} \quad (22)$$

To estimate the mean $\langle \text{GWI}_C \rangle$ for fossil fuel CO_2 emitted into the atmosphere between 1860 and 1989, I applied the exponential decay model (with $\tau_{\text{CO}_2} = 100 \pm 30$ yr) to the fossil fuel CO_2 emissions profile shown in Figure 1. This analysis yields $\langle \text{GWI}_C \rangle = 6.4 \pm 0.5 \mu\text{J m}^{-2}$ per kg C, essentially identical with the value obtained above with the exponential growth model, $6.2 \pm 0.5 \mu\text{J m}^{-2}$ per kg C. This result supports application of the heuristic exponential growth model given above to the interpretation of the relative forcings of CO_2 and sulfate over the industrial era. It may be noted that the value of $\langle \text{GWI}_C \rangle$ is relatively insensitive to the choice of τ_{CO_2} , an uncertainty in τ_{CO_2} of $\pm 30\%$ translating into an uncertainty in $\langle \text{GWI}_C \rangle$ of $\pm 5\%$. The reason for this is that most of the CO_2 has been emitted in the past 50 years, over which time GWI_C is quite insensitive to τ_{CO_2} ; cf. Figure 3. In addition to using the exponential decay profile for CO_2 , I also evaluated $\langle \text{GWI}_C \rangle$ using the nonexponential decay curve of Maier-Reimer and Hasselmann; {Maier-Reimer, 1987 #21} the resulting value, $5.7 \mu\text{J m}^{-2}$ per kg C, is nearly the same as the values obtained with the exponential decay curves. This agreement between the models shows that at least over the time period pertinent to this integration, estimation of the integrated CO_2 forcing is insensitive to the choice of model. The several estimates for GWI_C are summarized in Table 2.

Because the value of GWI_C obtained by integration over historical emissions is essentially the same as the value obtained with the model for exponentially increasing emissions, the estimate of $(Q_S / Q_C)^*$ is also the same as that obtained with that model. That is, the net forcing by CO_2 and sulfate over the industrial era are essentially identical, within the substantial uncertainty of the sulfate forcing. The several estimates for $(Q_S / Q_C)^*$ are summarized in Table 3.

Summary and Conclusions

A formalism involving Global Warming Integrals (GWI's), based on the Global Warming Potential concept, has been introduced to evaluate and compare the contributions to global warming and cooling from emissions of CO₂ and SO₂, respectively. The total GWI is appropriate for steady state emissions and for consideration of the total influence of CO₂ over its entire residence time in the atmosphere, but, in view of the rapid increase in the rate of CO₂ emissions on a time scale short to that residence time, this quantity substantially overestimates the influence that fossil fuel CO₂ has exerted on climate up to the present time. The partial influence up to the present time was evaluated by two models, an exponential growth model (e-folding time 30 years) and direct integration over historical emissions, yielding essentially the same results. The positive CO₂ forcing and the negative sulfate forcing were compared by comparing the critical sulfur to carbon mass ratio in emissions that would result in zero net forcing, $(Q_S/Q_C)^*$, with the actual ratio characterizing fossil fuel emissions over the industrial era $(\overline{Q_S}/\overline{Q_C})$. These comparisons indicate that the positive CO₂ forcing and the negative sulfate forcing, by direct light scattering only, are essentially equal, although the uncertainty in the latter is roughly a factor of two. This equality indicates that the net influence of fossil fuel effluents on radiative forcing of climate over the industrial period is near zero, consistent with the estimates of present net forcing, on which the calculations of GWI's were based. On the other hand the analysis shows that despite this near-zero present net forcing, the net influence over the entire projected residence time of the fossil fuel CO₂ in the atmosphere is positive by a substantial margin.

As noted above the GWI for sulfate, and therefore $(Q_S/Q_C)^*$, was evaluated for the forcing due only to direct light scattering, neglecting the cloud forcing. This cloud forcing has been estimated to be roughly equal globally to the direct forcing, but is considered less certain because of the untested assumptions that led to this estimate. Use of a value for GWI_S that includes both effects decreases estimates of $(Q_S/Q_C)^*$ by a further factor of about 2, i.e., to 0.038 for GWI_C^∞ and to 0.0085 for $\langle \text{GWI}_C \rangle$. The latter figure is well less than the historical average sulfur-to-carbon emission ratio (0.019), indicative, if it is accurate, that the net radiative forcing resulting from fossil fuel combustion over this period has been one of cooling, not warming. However this conclusion must remain tentative in view of the uncertainty in estimates of the radiative forcing of sulfate aerosol, {Wigley, 1992 #4; Charlson, 1992 #9} and the issue can be resolved only by decreasing this uncertainty.

Three final points should be noted. First, the calculations were carried out for a global-mean forcing, whereas in view of spatial and temporal differences in these forcings, substantial net short-term regional forcings of differing sign may be expected. Second, because of differing feedbacks in different regions, e.g., a greater water vapor feedback over tropical oceans, where the sulfate forcing is minimal, the mean global response to these disparate forcings might not be the same as that to a uniform forcing at the mean value. Third, in contrast to the sulfate forcing, the full effect of the CO₂ emissions has yet to be exerted. As noted previously{Wigley, 1991 #5; Kaufman, 1991 #6; Charlson, 1991 #7; Twomey, 1991 #8} inevitably SO₂ emissions must decrease, and subsequently, as the balance of the CO₂ forcing is exerted, this forcing must result in a much greater positive global-mean forcing than has been experienced thus far.

REFERENCES

- Charlson, R. J., Langner, J., Rodhe, H., Leovy, C. B. and Warren, S. G. *Tellus* **43AB**, 152 (1991);
 Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A. Jr., Hansen, J. E., and Hofmann, D. J. *Science* **255**, 423 (1992).
- Cullis, C. F. and Hirschler, M. M. *Atmos. Environ.* **14** 1263 (1980);
- Dignon, J. and Hameed, S. *JAPCA (J. Air Poll Contr. Assoc.)* **39** 180 (1989);
- Fisher, D. A., Hales, C. H., Wang, W.-C., Ko, M. K. W. and Sze, N. D. *Nature* **344** 513 (1990).
- Houghton, J. T., Jenkins, G. J. and Ephraums, J. J. *Climate Change: The IPCC Scientific Assessment*. (Cambridge University Press, Cambridge, 1990).
- Houghton, J.T., Callander, B.A. and Varney, S.K. *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment* (Cambridge Univ. Press, Cambridge, 1992).
- Isaksen et al in IPCC 1992
- Kaufman, Y. J., Fraser, R. S. and Mahoney, R. L. *J. Climate* **4**, 578 (1991);
- Kelly 1992
- Langner, J. and Rodhe H., *J. Atmos. Chem.* **13**, 225-263, 1991
- Maier-Reimer, E. and Hasselmann, K. *Climate Dynamics* **2**, 63 (1987).
- Marland, G., Boden, T. A., Griffin, R. C., Huang, S. F., Kanciruk, P. and Nelson, T. R. *Estimates of CO₂ Emissions from Fossil Fuel Burning and Cement Manufacturing, Based on the United Nations Energy Statistics and the U. S. Bureau of Mines Cement Manufacturing Data*. (Report ORNL / CDIAC-25, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1989).
- Möller, D. *Atmos. Environ.* **18** 19 (1984);
- Post, W. M., Peng, T.-H., Emanuel, W. R., King, A. W., Dale, V. H. and DeAngelis, D. L. *Amer. Scientist* **78** 310 (1990).
- Quay, P. D., Tilbrook, B. and Wong, C. S. *Science* **256** 74 (1992).
- Robinson, E. and Robbins, R. C. *Sources, Abundance and Fate of Gaseous Atmospheric Pollutants--Supplement*. (Report 4015, American Petroleum Institute, Washington DC, 1971);
- Schlesinger 1992
- Schwartz, S. E. *Nature* **336**, 441 (1988).
- Schwartz, S. E. *Tellus* **31**, 530 (1979).
- Shine 1990 in IPCC
- Siegenthaler, U. *J. Geophys. Res.* **88** 3599 (1983);

- Spiro, P.A., Jacob, D.J. and Logan, J.A. *J. Geophys. Res.* **97**, 6023 (1992).
Tans, P. P., Fung, I. Y. and Takahashi, T. *Science* **247** 1431 (1990);
Twomey, S. *Atmos. Environ.* **25A** 2435 (1991).
Twomey, S.A., Piepgrass, M. and Wolfe, T. L. *Tellus* **36B** 356 (1984).
Wang, W.-C., Dudek, M. P., Liang, X. Z., and Kiehl, J. T. *Nature* **350** 573 (1991).
Watson 1990 in IPCC
Wigley, T. M. L. *Nature* **349**, 503 (1991);
Wigley, T.M.L. and Raper, S.C.B. *Nature* **357**, 293 (1992).

ACKNOWLEDGMENT. This research was performed under the auspices of the United States Department of Energy, under Contract No. DE-AC02-76CH00016.

Model	----- Direct Forcing -----			Cloud Forcing
	Box ^(a)	3-D Global ^(b)	Employed in present evaluations ^(c)	Box ^(a)
$Q_S, \text{Tg S yr}^{-1}$	90	71	(81)	90
$Q_S, \text{kg S s}^{-1}$	2.85×10^3	2.25×10^3	(2.58×10^3)	2.85×10^3
$F_S, \text{W m}^{-2}$	-1.3	-0.6	(-1.0)	-1
$\text{GWI}_S^\infty, \mu\text{J m}^{-2} \text{kg}^{-1}$	-460	-270	-360	-360
$\tau_{\text{SO}_4^{2-}}, \text{days}$	7	5	6	7
$\tau_{\text{SO}_4^{2-}}, \text{s}$	6.05×10^5	4.32×10^5	5.18×10^5	6.05×10^5
$a_S, \text{pW m}^{-2} \text{kg}^{-1}$	-750	-620	-690	-580

Table 1. Comparison of GWI's for shortwave radiative forcing by sulfate aerosol evaluated with forcing calculated by box model and three-dimensional global model.

(a) Charlson et al.{Charlson, 1992 #9}

(b) Charlson et al.{ Charlson, 1991 #7}

(c) Entries in parentheses, which are not employed in present evaluations, are given to permit comparison.

	GWI_C^∞	$\text{GWI}_C^{\text{exp}}$ (30-year e-folding time of emissions)	$\langle \text{GWI}_C \rangle$ (1860-1989)
Decay Profile	----- $\mu\text{J m}^{-2}$ per kg C -----		
Exponential			
$\tau_{\text{CO}_2} = 70$ years	19.0	5.71	5.94
$\tau_{\text{CO}_2} = 100$ years	27.2	6.27	6.52
$\tau_{\text{CO}_2} = 130$ years	35.3	6.62	6.88
Maier-Reimer and Hasselmann{Maier -Reimer, 1987 #21}	-----	-----	5.71

Table 2. Comparison of GWI's for longwave radiative forcing by CO_2 for exponential and nonexponential models for the decay profile, f_{CO_2} , and for different approaches to the evaluation.

Critical Mass Emissions Ratio (Q_S / Q_C) [*]		
Model for (Q_S / Q_C) [*]	Expression	Value ^(a)
Steady state	$-GWIC^\infty / GWS^\infty$	0.075
Exponential growth ($\tau_e = 30$ years)	$-GWIC^{\text{exp}} / GWS^\infty$	0.017
Integrated (1860-1989)	$-\langle GWIC \rangle / GWS^\infty$	0.017
Actual Emissions Ratio (1860-1989)	$\overline{(Q_S / Q_C)}$	0.019

Table 3. Estimates of critical values of mass emissions ratio of sulfur to carbon from fossil fuel combustion (Q_S / Q_C)^{*} that would result in zero net radiative forcing of climate, according to different models of time-dependent emissions. Also shown is the average value $\overline{(Q_S / Q_C)}$ from fossil fuel combustion from 1860-1989. If the actual emissions ratio $\overline{(Q_S / Q_C)}$ exceeds (Q_S / Q_C)^{*}, then the net influence is cooling; if $\overline{(Q_S / Q_C)}$ is less than (Q_S / Q_C)^{*}, the net influence is warming. The uncertainty in (Q_S / Q_C)^{*} is estimated as factor of 2. The estimate of (Q_S / Q_C)^{*} does not include any contribution from enhanced cloud albedo, which would decrease its value.

(a) Estimates of (Q_S / Q_C)^{*} are uncertain to a factor of 2, largely due to uncertainty in the estimate of GWS^∞ .




Figure 1. Time history of ratio of sulfur to carbon emissions from fossil fuel combustion evaluated from Refs.^{16, 24-27} (left axis) and the rate of CO₂ emissions from fossil fuel combustion²⁸ (right axis).




Figure 2. Profiles of fraction of excess CO₂ remaining in atmosphere as function of time subsequent to introduction into atmosphere. Solid curves show exponential decay for indicated mean residence times in years. Dotted curve shows nonexponential decay curve obtained with multi-component box model, as employed in Global Warming Potential estimates by IPCC.¹⁴




Figure 3. Profiles of the time integral of the fraction of excess CO_2 remaining in atmosphere as function of time subsequent to introduction into atmosphere. Solid curves shows integral of exponential decay for indicated mean residence times in years. Dotted curve shows integral of nonexponential decay curve obtained with multi-component box model. Right hand axis shows Global Warming Integral evaluated for the instantaneous radiative forcing due to an incremental amount of CO_2 in the atmosphere $a_{\text{CO}_2} = 8.6 \times 10^{-15} \text{ W m}^{-2} \text{ per kg C}$.




Figure 4. Rate of anthropogenic CO₂ emissions from fossil fuel combustion over the industrial era.²⁸ Straight line represents exponential growth with 30 year e-folding time.